



PATENT APPLICATION

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of

Nobuo HANAI *et al.*

Atty. Ref.: 249-255

Appln. No.: 10/089,500

Group Art Unit: 1642

Filed: March 29, 2002

Examiner: Blanchard, David J

For: COMPLEMENTARITY DETERMINING REGION-GRAFTED
ANTIBODY AGAINST GANGLIOSIDE GD3 AND
DERIVATIVE OF ANTIBODY AGAINST GANGLIOSIDE GD3

DECLARATION

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir/Madam:

I, Eiichi Kobayashi, do declare and state that:

I graduated from the University of Tokyo, Faculty of Agriculture, Department in Agricultural Chemistry, having received a Master's Degree of Agriculture in March, 1992.

I understand the Japanese and English languages.

I understand the Japanese and English languages. Attachment is an accurate English translation made by me of Japanese Patent Application No. 2000-105088 filed on April 6, 2000.

I declare further that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Date: July 13, 2005

Name:

Eiichi Kobayashi

Best Available Copy

[Document]	Patent Application
[Reference No.]	H11-2101Q3
[Date of filing]	April 6, 2000
[Attention]	Commissioner, Patent Office
[ITC]	C12P 21/08
[Inventor]	
[Address]	c/o Tokyo Research Laboratories, Kyowa Hakko Kogyo Co., Ltd., 6-6, Asahi-machi 3-chome, Machida-shi, Tokyo, Japan
[Name]	Nobuo HANAI
[Inventor]	
[Address]	c/o Tokyo Research Laboratories, Kyowa Hakko Kogyo Co., Ltd., 6-6, Asahi-machi 3-chome, Machida-shi, Tokyo, Japan
[Name]	Kenya SHITARA
[Inventor]	
[Address]	c/o Tokyo Research Laboratories, Kyowa Hakko Kogyo Co., Ltd., 6-6, Asahi-machi 3-chome, Machida-shi, Tokyo, Japan
[Name]	Kazuyasu NAKAMURA
[Inventor]	
[Address]	c/o Tokyo Research Laboratories, Kyowa Hakko Kogyo Co., Ltd., 6-6, Asahi-machi 3-chome, Machida-shi, Tokyo, Japan
[Name]	Rinpei NIWA
[Applicant for patent]	
[Identification No.]	000001029
[Name or appellation]	Kyowa Hakko Kogyo Co., Ltd.
[Representative]	Tadashi HIRATA
[Domestic priority to be claimed based on the prior application]	
[Application No.]	Hei. 11-278291
[Filing date]	September 30, 1999
[Identification of fees]	
[Prepayment account No.]	008187
[Amount of fees payable]	21000
[List of documents to be submitted]	
[Document]	Specification 1
[Document]	Drawing 1
[Document]	Abstract 1
[Request of Proof]	Yes

(Document name)

Specification

(Title of the invention)

COMPLEMENTARITY DETERMINING REGION-GRAFTED
ANTIBODY AGAINST GANGLIOSIDE GD3 AND
DERIVATIVE OF ANTIBODY AGAINST GANGLIOSIDE GD3

(Scope of the claims)

(Claim 1) A derivative of an antibody, comprising a monoclonal antibody or the antibody fragment thereof which specifically reacts with ganglioside GD3 and is conjugated with a radioisotope, a protein or a low molecular weight agent.

(Claim 2) The derivative of an antibody according to claim 1, wherein the monoclonal antibody which specifically reacts with ganglioside GD3 is an antibody selected from an antibody produced by a hybridoma, a humanized antibody and a human antibody.

(Claim 3) The derivative of an antibody according to claim 2, wherein the antibody produced by a hybridoma is KM641.

(Claim 4) The derivative of an antibody according to claim 2, wherein the humanized antibody is a human chimeric antibody or a human CDR-grafted antibody.

(Claim 5) The derivative of a human chimeric antibody according to claim 4, wherein the human chimeric antibody comprises an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3 produced by a hybridoma.

(Claim 6) The derivative of a human chimeric antibody according to claim 4, wherein the human chimeric antibody comprises: an H chain V region and an L chain V region of a monoclonal antibody produced against ganglioside GD3 by a hybridoma; and an H chain constant region (C region) and an L chain C region of a human antibody.

(Claim 7) The derivative of a human chimeric antibody according to claim 5 or 6, wherein the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55.

(Claim 8) The derivative of a human chimeric antibody according to claim 5 or 6, wherein the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

(Claim 9) The derivative of a human chimeric antibody according to claim 5 or 6, wherein the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55; and the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

(Claim 10) The derivative of a human chimeric antibody KM871 according to claim 5 or 6, wherein the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55; and the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

(Claim 11) The derivative of a human CDR-grafted antibody according to claim 4, wherein the human CDR-grafted antibody comprises CDR of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3.

(Claim 12) The derivative of a human CDR-grafted antibody according to claim 4, wherein the human CDR-grafted antibody comprises: CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3; and framework regions (FRs) of an H chain V region and an L chain V region of a human antibody.

(Claim 13) The derivative of an antibody according to claim 47, wherein the human CDR-grafted antibody comprises: CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3; FRs of an H chain V region and an L chain V region of a human antibody; and an H chain C region and an L chain C region of a human antibody.

(Claim 14) The derivative of a human CDR-grafted antibody according to any one of claims 11 to 13, wherein the antibody comprises CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

(Claim 15) The derivative of a human CDR-grafted antibody according to any one of claims 11 to 13, wherein the antibody comprises CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

(Claim 16) The derivative of a human CDR-grafted antibody according to any one of claims 11 to 13, wherein the antibody comprises: CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively; and CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

(Claim 17) The derivative of a human CDR-grafted antibody according to any one of claims 11 to 13, wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9.

(Claim 18) The derivative of a human CDR-grafted antibody according to any one of claims 11 to 13, wherein the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

(Claim 19) The derivative of a human CDR-grafted antibody according to any one of claims 11 to 13, wherein the H chain V region and the L chain V region of the antibody comprises the amino acid sequences represented by SEQ ID NO:9 and SEQ ID NO:54, respectively.

(Claim 20) The derivative of a human CDR-grafted antibody KM8871 according to any one of claims 11 to 13, wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

(Claim 21) The derivative of the antibody fragment according to claim 1, wherein the antibody fragment is an antibody fragment selected from Fab, Fab', F(ab')₂, a single chain antibody (scFv), a disulfide stabilized V region fragment (dsFv) and a peptide comprising CDR.

(Claim 22) The derivative of the antibody fragment according to claim 1 or 21, wherein the antibody fragment comprises amino acid sequences of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3 produced by a hybridoma.

(Claim 23) The derivative of the antibody fragment according to claim 1 or 21, wherein the antibody fragment comprises an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:55.

(Claim 24) The derivative of the antibody fragment according to claim 1 or 21, wherein the antibody fragment comprises an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:56.

(Claim 25) The derivative of the antibody fragment according to claim 1 or 21, wherein the antibody fragment comprises: an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:55; and an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:56.

(Claim 26) The derivative of the antibody fragment according to claim 1 or 21, wherein the antibody fragment comprises amino acid sequences of an H chain V region and an L chain V region of a human CDR-grafted antibody against ganglioside GD3.

(Claim 27) The derivative of the antibody fragment according to claim 1 or 21, wherein the antibody fragment comprises an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:9.

(Claim 28) The derivative of the antibody fragment according to claim 1 or 21, wherein the antibody fragment comprises an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:54.

(Claim 29) The derivative of the antibody fragment according to claim 1 or 21, wherein the antibody fragment comprises: an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:9; and an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:54.

(Claim 30) The derivative of an antibody fragment according to claim 1 or 21, wherein the antibody fragment comprises CDR1, CDR2 and CDR3 of an H chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

(Claim 31) The derivative of the antibody fragment according to claim 1 or 21, wherein the antibody fragment comprises CDR1, CDR2 and CDR3 of an L chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

(Claim 32) The derivative of the antibody fragment according to claim 1 or 21, wherein the antibody fragment comprises: CDR1, CDR2 and CDR3 of an H chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5; and CDR1, CDR2 and CDR3 of an L chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8.

(Claim 33) The derivative of a monoclonal antibody or the antibody fragment thereof according to any one of claim 1 to 32, wherein the protein is a cytokine.

(Claim 34) The derivative of a monoclonal antibody or the antibody fragment thereof according to claim 31, wherein the cytokine is human interleukin-2 (hIL-2).

(Claim 35) The derivative of an antibody according to claim 34, wherein the derivative of an antibody comprises a human chimeric antibody KM871 and hIL-2.

(Claim 36) The derivative of an antibody according to claim 35, wherein the antibody conjugated with hIL-2 comprises: an H chain V region having the amino acid sequence represented by SEQ ID NO:57; and an L chain V region having the amino acid sequence represented by SEQ ID NO:56.

(Claim 37) The derivative of an antibody according to claim 36, wherein the derivative of an antibody comprises a human CDR-grafted antibody KM8871 and hIL-2.

(Claim 38) The derivative of an antibody according to claim 1, wherein the antibody conjugated with hIL-2 comprises: an H chain V region having the amino acid sequence represented by SEQ ID NO:53; and an L chain V region having the amino acid sequence represented by SEQ ID NO:54.

(Claim 39) A DNA which encodes the derivative of a monoclonal antibody or the derivative of the antibody fragment thereof which specifically reacts with ganglioside GD3 according to any one of claims 1 to 38.

(Claim 40) A recombinant vector comprising the DNA according to claim 39.

(Claim 41) A transformant which is obtained by introducing the recombinant vector according to claim 40 into a host cell.

(Claim 42) A transformant KM871hIL2 (FERM BP-6918) which produces the antibody according to claim 35.

(Claim 43) A transformant KM8871hIL2 (FERM BP-6791) which produces the antibody according to claim 37.

(Claim 44) A process for producing an antibody, which comprises: culturing the transformant according to any one of claims 41 to 43 in a culture medium to produce and accumulate the derivative of a monoclonal antibody or the derivative of the antibody fragment thereof according to any one of claims 1 to 38 in the culture; and recovering the derivative of the antibody or the derivative of the antibody fragment thereof from the culture.

(Claim 45) A human CDR-grafted antibody or the antibody fragment thereof which specifically reacts with ganglioside GD3.

(Claim 46) The human CDR-grafted antibody or the antibody fragment thereof according to claim 45, wherein the human CDR-grafted antibody comprises CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3.

(Claim 47) The human CDR-grafted antibody or the antibody fragment thereof according to claim 45, wherein the human CDR-grafted antibody comprises: CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3; and FRs of an H chain V region and an L chain V region of a human antibody.

(Claim 48) The human CDR-grafted antibody or the antibody fragment thereof according to claim 45, wherein the human CDR-grafted antibody comprises: CDRs of an H chain V region and an L chain V region of a monoclonal antibody against

ganglioside GD3; FRs of an H chain V region and an L chain V region of a human antibody; and an H chain C region and an L chain C region of a human antibody.

(Claim 49) The human CDR-grafted antibody or the antibody fragment thereof according to any one of claims 46 to 48, wherein the antibody comprises CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

(Claim 50) The human CDR-grafted antibody or the antibody fragment thereof according to any one of claims 46 to 48, wherein the antibody comprises CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

(Claim 51) The human CDR-grafted antibody or the antibody fragment thereof according to any one of claims 46 to 48, wherein the antibody comprises: CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively; and CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8.

(Claim 52) The human CDR-grafted antibody or the antibody fragment thereof according to any one of claims 46 to 48, wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9.

(Claim 53) The human CDR-grafted antibody or the antibody fragment thereof according to any one of claims 46 to 48, wherein the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

(Claim 54) The human CDR-grafted antibody or the antibody fragment thereof according to any one of claims 46 to 48, wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

(Claim 55) The human CDR-grafted antibody KM8871 or the antibody fragment thereof according to any one of claims 46 to 48, wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

(Claim 56) A DNA which encodes the human CDR-grafted antibody or the antibody fragment thereof which specifically reacts with ganglioside GD3 according to any one of claims 45 to 55.

(Claim 57) A recombinant vector comprising the DNA according to claim 56.

(Claim 58) A transformant which is obtained by introducing the recombinant vector according to claim 57 into a host cell.

(Claim 59) A transformant KM8871 (FERM BP-6790) which produces the human CDR-grafted antibody according to claim 55.

(Claim 60) A process for producing an antibody, which comprises: culturing the transformant according to claim 58 or 59 in a culture medium to produce and accumulate the human CDR-grafted antibody or the antibody fragment thereof according to any one of claims 45 to 55 in the culture; and recovering the antibody or the antibody fragment thereof from the culture.

(Claim 61) A therapeutic agent for cancers, comprising, as an active ingredient, at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to any one of claims 1 to 38 and the human CDR-grafted antibody and the antibody fragment thereof according to any one of claims 45 to 55.

(Claim 62) A diagnostic agent for cancers, comprising, as an active ingredient, at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to any one of claims 1 to 38 and the human CDR-grafted antibody and the antibody fragment thereof according to any one of claims 45 to 55.

(Detailed description of the invention)

(0001)

(Technical field to which the invention belongs)

5 The present invention relates to derivatives of a monoclonal antibody and the antibody fragment which specifically react with ganglioside GD3 (hereinafter referred to as "GD3"), and to a human CDR-grafted antibody and the antibody fragment which specifically react with GD3. Furthermore, the present invention relates to a DNA sequence which encodes the derivatives, antibody or antibody fragment. The present invention relates to a vector comprising the DNA sequence and a cell
10 transformed with the vector. Moreover, the present invention relates to a process for producing the derivatives, antibody and antibody fragment using the transformed cell, and to a diagnostic agent and a therapeutic agent for cancers using the derivatives, antibody and antibody fragment.

(0002)

15 (Background art)

It is known that, when an antibody derived from a non-human animal, e.g., a mouse antibody, is administered to human, it is recognized as a foreign substance and

thus induces a human antibody against the mouse antibody in the human body (human anti mouse antibody, hereinafter referred to as "HAMA"). It is known that the HAMA reacts with the administered mouse antibody to cause side effects (*J. Clin. Oncol.*, 2, 881 (1984); *Blood*, 65, 1349 (1985); *J. Natl. Cancer Inst.*, 80, 932 (1988); *Proc. Natl. Acad. Sci. U.S.A.*, 82, 1242 (1985)), to quicken disappearance of the administered mouse antibody from blood (*J. Nucl. Med.*, 26, 1011 (1985); *Blood*, 65, 1349 (1985); *J. Natl. Cancer Inst.*, 80, 937 (1988)) and to reduce therapeutic effects of the mouse antibody (*J. Immunol.*, 135, 1530 (1985); *Cancer Res.*, 46, 6489 (1986)).

(0003)

In order to resolve these problems, attempts have been made to genetic engineeringly convert an antibody derived from a non-human animal into a humanized antibody such as a human chimeric antibody or a human complementarity determining region (hereinafter referred to as "CDR")-grafted antibody. The human chimeric antibody is an antibody in which its antibody variable region (hereinafter referred to as "V region") is derived from a non-human animal antibody and its constant region (hereinafter referred to as "C region") is derived from a human antibody (*Proc. Natl. Acad. Sci. U.S.A.*, 81, 6851 (1984)). The human CDR-grafted antibody is an antibody in which the amino acid sequence of CDR in the V region derived from a non-human animal antibody is grafted to an appropriate position of a human antibody (*Nature*, 321, 522 (1986)). These humanized antibodies have various advantages in clinically applying to human, in comparison with antibodies of non-human animals such as mouse antibodies and the like. For example, it has been reported on the immunogenicity and stability in blood that, when a human chimeric antibody is administered to human, its half-life in blood increased about six times in comparison with a mouse antibody (*Proc. Natl. Acad. Sci. U.S.A.*, 86, 4220 (1989)). It has been reported that, in a test of a human CDR-grafted antibody in monkeys, its immunogenicity is reduced and its half-life in blood is increased four to five times in comparison with a mouse antibody (*J. Immunol.*, 147, 1352 (1991)). That is, it is expected that humanized antibodies have less side effects and their therapeutic effects continue for a longer period of time than antibodies of non-human animals. In addition, when applications particularly as anti-tumor antibodies are taken into consideration, higher cytotoxic activities such as complement-dependent cytotoxicity (hereinafter referred to as "CDC"), antibody-dependent cell-mediated cytotoxic activity (hereinafter referred to as "ADCC") and the like via an antibody Fc region (region in and after the hinge region of an antibody heavy chain) are important for the therapeutic effects. It has been reported that on such cytotoxic activities, that the Fc region of human antibodies is more superior in human to

the Fc region of antibodies of non-human animals, since the Fc region more effectively activates human complement components and human effector cells having Fc receptors of mononuclear cells, macrophages and NK cells on the cell surface. For example, it has been reported that when a mouse antibody against GD3 (hereinafter referred to as "anti-GD3 mouse antibody") is converted into a human chimeric antibody having a human antibody Fc region (hereinafter referred to as "anti-GD3 chimeric antibody"), its tumor cell growth-inhibitory activity by human effector cells is increased (*J. Immunol.*, 144, 1382 (1990)), and similar results have been reported on a human CDR-grafted antibody against CAMPATH-1 antigen (*Nature*, 332, 323 (1988)).

(0004)

These results clearly show that humanized antibodies are preferred to non-human antibodies such as mouse antibodies and the like, as antibodies for clinical applications to human.

In addition, with the recent advance in protein engineering and genetic engineering, antibody fragments having a smaller molecular weight such as Fab, Fab', F(ab')₂, a single chain antibody (*Science*, 242, 423 (1988)), a disulfide stabilized V region fragment (*Molecular Immunol.*, 32, 249 (1995)) and the like, can be produced. Since these fragments have a smaller molecular weight than whole antibody molecules, they are excellent in penetrating ability into target tissues (*Cancer Res.*, 52, 3402 (1992)). It is considered that these fragments derived from humanized antibodies are more desirable than those derived from non-human animal antibodies such as mouse antibodies, when used in clinical applications to human.

(0005)

Ganglioside which is one of glycolipids containing sialic acid is a molecule which constitutes animal cell membrane and is constituted by a sugar chain as a hydrophilic side chain and a sphingosine and fatty acid as hydrophobic side chains. It is known that kinds and expression levels of ganglioside vary depending on the cell types, organ species, animal species and the like. It is also known that the expression of ganglioside changes quantitatively and qualitatively in the process of malignant transformation of cells (*Cancer Res.*, 45, 2405 (1985)). Particularly, the GD3 relating to the present invention is present in an extremely small amount in normal cells but in a large amount in cancer cells such as melanoma, sarcoma, glioma, neuroblastoma and the like (*Proc. Natl. Acad. Sci. U.S.A.*, 77, 6114 (1980); *J. Exp. Med.*, 155, 1133 (1982); *Cancer Res.*, 45, 4401 (1985)), and an antibody against GD3 (hereinafter referred to as "anti-GD3 antibody") is considered to be useful for the treatment of these cancers (*Melanoma Research*, 7, S115 (1997)). Tests on its administration to melanoma

patients have already been carried out using R24 as an anti-GD3 mouse antibody, and its clinical effect was observed in a part of the patients but not to an expected degree because of considerably quick disappearance of the antibody from blood due to induction of HAMA (*Melanoma Research*, 7, S115 (1997)). Under such
5 circumstances, in order to achieve more effective clinical applications of an anti-GD3 antibody to human, the present inventors have attempted to produce an anti-GD3 chimeric antibody having similar activity to that of a mouse antibody based on an anti-GD3 mouse antibody KM641 produced by the inventors, and have succeeded in producing an anti-GD3 chimeric antibody KM871 (Japanese Published Unexamined
10 Patent Application No. 304989/93). It is expected that, in comparison with the anti-GD3 mouse antibody, the anti-GD3 chimeric antibody KM871 reduces immunogenicity in human, prolongs half-life in blood and has a stronger anti-tumor effect. However, it was found as a result of clinical tests on other anti-GD3 chimeric antibodies in human that allergy reactions are induced due to induction of a human antibody against a V
15 region derived from a mouse antibody in the anti-GD3 chimeric antibodies (*Cancer J. From Scientific American*, 3, S121 (1997)). Even in the anti-GD3 chimeric antibody KM871 prepared by the inventors, it is necessary to await results of clinical tests in human to know if similar reactions are induced or not, and in order to reduce a possibility of generating such a problem as low as possible, it is desirable to produce a
20 human CDR-grafted antibody against GD3 (hereinafter referred to as "anti-GD3 CDR-grafted antibody") which is considered to have lower immunogenicity than those of anti-GD3 chimeric antibodies in human. However, to date, there are no reports on the production of an anti-GD3 CDR-grafted antibody.

(0006)

25 With regard to antibody fragments, a single chain antibody derived from an anti-GD2 mouse antibody has been produced (*Hybridoma*, 16, 335 (1997)), but there are no reports on the production of other antibody fragments. With regard to antibody fragments derived from an anti-GD3 antibody, there are no reports on the production of antibody fragments derived from a mouse antibody and a humanized antibody. Thus,
30 if an antibody fragment derived from a humanized antibody against GD3 (hereinafter referred to as "anti-GD3 humanized antibody") can be produced, it is expected that it has excellent target tissue penetratability and the immunogenicity is reduced.

(0007)

35 As has been described above, it is expected that a humanized antibody and the antibody fragment have effect on diagnostic and therapeutic methods by single use, but further improvement of the effects has been examined by using them in combination

with other molecules. For example, cytokine is used as one of these molecules. Cytokine is a general term for various aqueous factors which control intercellular mutual reactions in immune interactions. As cytotoxic activities of antibodies such as CDC activity, ADCC activity and the like are known and the ADCC activity is effected
5 by effector cells having Fc receptors on the cell surface such as mononuclear cells, macrophages, NK cells and the like (*J. Immunol.*, 138, 1992 (1987)). Since various kinds of cytokine activate these effector cells, their administration in combination with antibodies has been carried out to improve ADCC activity and the like of antibodies. For example, with regard to an anti-GD2 mouse antibody 14.G2a and an anti-GD2
10 chimeric antibody ch14.18, their administration to human in combination with a cytokine human interleukin-2 (hereinafter referred to as "hIL-2") or a human granulocyte macrophage colony-stimulating factor (hereinafter referred to as "hGM-CSF") has been carried out (*Cancer*, 80, 317 (1997); *Cancer J. From Scientific American*, 3, S121 (1997)). Regarding an anti-GD3 mouse antibody R24, combination
15 therapy with various kinds of cytokine has also been carried out (*Cancer Res.*, 50, 7490 (1990), *Proc. Am. Soc. Clin. Oncol.*, 1186, 345 (1992), *J. Biol. Response Mod.*, 9, 319 (1990), *Proc. Am. Soc. Clin. Oncol.*, 1182, 344 (1992), *Proc. Am. Soc. Clin. Oncol.*, 1188, 346 (1992)). However, the effects as to be expected are not found in these combination therapies due to immunogenicity of mouse antibodies or side effects
20 derived from cytokine. Thus, derivatives has been produced by conjugating an antibody or the antibody fragment thereof with a radioisotope, a protein (cytokine, toxin, enzyme or the like), a low molecular weight agent and the like, chemically or genetic engineeringly, and their clinical applications have been examined (*New Eng. J. Med.*, 329, 459 (1993), *Anticancer Res.*, 17, 1735 (1997), *Blood*, 78, 1173 (1991), *J. Clin. Oncol.*, 15, 723 (1997), *Bioconjugate Chem.*, 7, 606 (1997), *Cancer*, 61, 881 (1988),
25 *Jpn. J. Cancer Res.*, 85, 167 (1994), *Antibody Immunoconjugates and Radiopharmaceuticals*, 3, 60 (1990), *Surgery*, 106, 533 (1989)). It is expected that these derivatives can accumulate a radioisotope, a protein (cytokine, toxin, enzyme or the like), a low molecular weight agent or the like in the periphery of a target tissue according to the binding specificity of antibodies so that more effective diagnoses and
30 treatments having less side effects can be obtained. For example, with regard to hIL-2 which showed a certain degree of anti-tumor effect among cytokines used in the above combination therapy, its fusion protein with an anti-GD2 chimeric antibody ch14.18 was produced genetic engineeringly, and it has been reported that its antitumor effects
35 in tests using mice were superior to those of simultaneous administration of anti-GD2 chimeric antibody ch14.18 and hIL-2 (*Proc. Natl. Acad. Sci. U.S.A.*, 89, 1428 (1992),

Proc. Natl. Acad. Sci. U.S.A., 91, 9626 (1994), *Cancer Immunol. Immunother.*, 42, 88 (1996), *Blood*, 91, 1706 (1998)). However, with regard to an anti-GD3 antibody, there are no reports on the production of derivatives by conjugating it with a protein (cytokine, toxin, enzyme or the like), a radioisotope, a low molecular weight agent and the like.

5 Accordingly, also in the case of the anti-GD3 humanized antibody and the antibody fragment of the present invention, if derivatives conjugated with a radioisotope, a protein (toxin, enzyme or the like), a low molecular weight agent and the like, including fusion proteins with various kinds of cytokine, can be produced, reduction of immunogenicity, less side effects and more potent antitumor effects in the tumor region
10 can be expected when administered to the human body.

(0008)

(Problems to be solved by the invention)

An object of the present invention is to provide an anti-GD3CDR-grafted antibody or the antibody fragment which has lower immunogenicity in human than an
15 anti-GD3 mouse antibody and an anti-GD3 chimeric antibody and which has potent anti-tumor effect. Also, an object of the present invention is to produce derivatives of an antibody or an antibody fragment thereof which is a fusion protein, said fusion protein comprises an anti-GD3 monoclonal antibody, especially an anti-GD3 humanized antibody or the antibody fragment thereof such as an anti-GD3 chimeric antibody or
20 anti-GD3CDR-grafted antibody, fused with cytokine, *etc.* and to accomplish therapeutic effects such as potent local anti-tumor effect at human tumors using the derivatives.

(0009)

(Means to solve the problems)

The inventors of the present invention obtained cDNAs encoding antibody
25 heavy chain (hereinafter referred to as H chain) and light chain (hereinafter referred to as L chain) from hybridoma cell KM641 which produces an anti-GD3 mouse antibody KM641 belonging to the IgG3 class, constructed an anti-GD3 chimeric antibody expression vector by cloning cDNAs encoding the V regions thereof into an expression vector for animal cell having cDNAs encoding human antibody H chain and L chain C regions, and prepared a transformant KM871 (FERM BP-3512) which produces an anti-
30 GD3 chimeric antibody KM871 in culture supernatant by introducing the expression vector into an animal cell. Furthermore, they purified an anti-GD3 chimeric antibody KM871 from the supernatant of the transformant KM871 and clarified that the anti-GD3 chimeric antibody KM871 has antigen binding activity, antigen binding specificity and
35 potent anti-tumor effect against a human cancer clone-grafted mouse, each activity being similar to that of the anti-GD3 mouse antibody KM641, so that demonstrate

utility of the anti-GD3 chimeric antibody KM871 in clinical applications to human (Japanese Published Unexamined Patent Application No. 304989/93).

(0010)

In the present invention, an amino acid sequence of each CDR was identified from the amino acid sequences of H chain and L chain V regions of the anti-GD3 mouse antibody KM641, cDNAs encoding the H chain and L chain V regions comprising the amino acid sequences of the CDRs of KM641 and amino acid sequences of the framework regions (hereinafter referred to as "FR") of human antibody H chain and L chain V regions were cloned into an expression vector for animal cell having cDNA encoding human antibody H chain and L chain C regions, and then the thus constructed anti-GD3 CDR-grafted antibody expression vector was introduced into an animal cell to produce a transformant KM8871 (FERM BP-6790) which produces an anti-GD3 CDR-grafted antibody KM8871 in a culture supernatant. Next, the anti-GD3 CDR-grafted antibody KM8871 was purified from a culture supernatant of the transformant KM8871 to find that the anti-GD3 CDR-grafted antibody KM8871 shows an antigen binding activity, an antigen binding specificity and a strong cytotoxic activity against human cancer cell lines, which are similar to those of the anti-GD3 chimeric antibody KM871.

(0011)

Furthermore, a cDNA was constructed by linking other cDNA encoding hIL-2 to the 3'-terminal of cDNA encoding the H chain of an anti-GD3 antibody, particularly anti-GD3 CDR-grafted antibody KM8871, and then the cDNA and a cDNA encoding the L chain of KM8871 were cloned into an expression vector for animal cell to construct an expression vector of a fusion protein of anti-GD3 CDR-grafted antibody KM8871 with hIL-2 (hereinafter referred to as "KM8871-hIL-2"). A transformant KM8871hIL2 (FERM BP-6790) which produces KM8871-hIL-2 in culture supernatant was produced by introducing the KM8871-hIL-2 expression vector into an animal cell. Furthermore, KM8871-hIL-2 was purified from a culture supernatant of the transformant KM8871hIL2 to find that KM8871-hIL-2 shows an antigen binding activity and an antigen binding specificity, which are similar to those of anti-GD3 CDR-grafted antibody KM8871, and a growth supporting activity against cell lines showing hIL-2-dependent growth, similar to that of hIL-2. Thereafter, the present invention was accomplished by finding that an activity of KM8871-hIL-2 in terms of the cytotoxic activity measured using a human peripheral blood mononuclear cell fraction is improved in comparison with that of anti-GD3 CDR-grafted antibody KM8871.

(0012)

The present invention relates to the following (1) to (62).

(1) A derivative of an antibody, comprising a monoclonal antibody or the antibody fragment thereof which specifically reacts with ganglioside GD3 and is conjugated with a radioisotope, a protein or a low molecular weight agent.

5 (2) The derivative of an antibody according to (1), wherein the monoclonal antibody which specifically reacts with ganglioside GD3 is an antibody selected from an antibody produced by a hybridoma, a humanized antibody and a human antibody.

(0013)

10 (3) The derivative of an antibody according to (2), wherein the antibody produced by a hybridoma is KM641 (FERM BP-3116).

(4) The derivative of an antibody according to (2), wherein the humanized antibody is a human chimeric antibody or a human CDR-grafted antibody.

15 (5) The derivative of a human chimeric antibody according to (4), wherein the human chimeric antibody comprises an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3 produced by a hybridoma.

(0014)

20 (6) The derivative of a human chimeric antibody according to (4), wherein the human chimeric antibody comprises: an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3 produced by a hybridoma; and an H chain constant region (C region) and an L chain C region of a human antibody.

(7) The derivative of a human chimeric antibody according to (5) or (6), wherein the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55.

(0015)

25 (8) The derivative of a human chimeric antibody according to (5) or (6), wherein the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

30 (9) The derivative of a human chimeric antibody according to (5) or (6), wherein the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55; and the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

(0016)

35 (10) The derivative of a human chimeric antibody KM871 according to (5) or (6), wherein the H chain V region comprises the amino acid sequence represented by SEQ ID NO:55; and the L chain V region comprises the amino acid sequence represented by SEQ ID NO:56.

(11) The derivative of a human CDR-grafted antibody according to (4), wherein the human CDR-grafted antibody comprises CDR of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3.

(0017)

5 (12) The derivative of a human CDR-grafted antibody according to (4), wherein the human CDR-grafted antibody comprises: CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3; and framework regions (FRs) of an H chain V region and an L chain V region of a human antibody.

10 (13) The derivative of an antibody according to (4), wherein the human CDR-grafted antibody comprises: CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3; FRs of an H chain V region and an L chain V region of a human antibody; and an H chain C region and an L chain C region of a human antibody.

(0018)

15 (14) The derivative of a human CDR-grafted antibody according to any one of (11) to (13), wherein the antibody comprises CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

20 (15) The derivative of a human CDR-grafted antibody according to any one of (11) to (13), wherein the antibody comprises CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

(0019)

25 (16) The derivative of a human CDR-grafted antibody according to any one of (11) to (13), wherein the antibody comprises: CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively; and CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

30 (17) The derivative of a human CDR-grafted antibody according to any one of (11) to (13), wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9.

(0020)

35 (18) The derivative of a human CDR-grafted antibody according to any one of (11) to (13), wherein the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

(19) The derivative of a human CDR-grafted antibody according to any one of (11) to (13), wherein the H chain V region and the L chain V region of the antibody comprises the amino acid sequences represented by SEQ ID NO:9 and SEQ ID NO:54, respectively.

5 (0021)

(20) The derivative of a human CDR-grafted antibody KM8871 according to any one of (11) to (13), wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

10 (21) The derivative of the antibody fragment according to (1), wherein the antibody fragment is an antibody fragment selected from Fab, Fab', F(ab')₂, a single chain antibody (scFv), a disulfide stabilized V region fragment (dsFv) and a peptide comprising CDR.

(0022)

15 (22) The derivative of the antibody fragment according to (1) or (21), wherein the antibody fragment comprises amino acid sequences of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3 produced by a hybridoma.

20 (23) The derivative of the antibody fragment according to (1) or (21), wherein the antibody fragment comprises an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:55.

(0023)

25 (24) The derivative of the antibody fragment according to (1) or (21), wherein the antibody fragment comprises an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:56.

(25) The derivative of the antibody fragment according to (1) or (21), wherein the antibody fragment comprises: an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:55; and an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:56.

30 (0024)

(26) The derivative of the antibody fragment according to (1) or (21), wherein the antibody fragment comprises amino acid sequences of an H chain V region and an L chain V region of a human CDR-grafted antibody against ganglioside GD3.

35 (27) The derivative of the antibody fragment according to (1) or (21), wherein the antibody fragment comprises an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:9.

(0025)

(28) The derivative of the antibody fragment according to (1) or (21), wherein the antibody fragment comprises an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:54.

5 (29) The derivative of the antibody fragment according to (1) or (21), wherein the antibody fragment comprises: an H chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:9; and an L chain V region of the antibody having the amino acid sequence represented by SEQ ID NO:54.

(0026)

10 (30) The derivative of an antibody fragment according to (1) or (21), wherein the antibody fragment comprises CDR1, CDR2 and CDR3 of an H chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

15 (31) The derivative of the antibody fragment according to (1) or (21), wherein the antibody fragment comprises CDR1, CDR2 and CDR3 of an L chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

(0027)

20 (32) The derivative of the antibody fragment according to (1) or (21), wherein the antibody fragment comprises: CDR1, CDR2 and CDR3 of an H chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5; and CDR1, CDR2 and CDR3 of an L chain V region of the antibody having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8.

25 (33) The derivative of a monoclonal antibody or the antibody fragment thereof according to any one of (1) to (32), wherein the protein is a cytokine.

(0028)

(34) The derivative of a monoclonal antibody or the antibody fragment thereof according to (33), wherein the cytokine is human interleukin-2 (hIL-2).

30 (35) The derivative of an antibody according to claim 34, wherein the derivative of an antibody comprises a human chimeric antibody KM871 and hIL-2.

(36) The derivative of an antibody according to (35), wherein the antibody conjugated with hIL-2 comprises: an H chain V region having the amino acid sequence represented by SEQ ID NO:57; and an L chain V region having the amino acid sequence represented by SEQ ID NO:56.

35 (0029)

(37) The derivative of an antibody according to (34), wherein the derivative of an antibody comprises a human CDR-grafted antibody KM8871 and hIL-2.

(38) The derivative of an antibody according to (1), wherein the antibody conjugated with hIL-2 comprises: an H chain V region having the amino acid sequence represented by SEQ ID NO:53; and an L chain V region having the amino acid sequence represented by SEQ ID NO:54.

(0030)

(39) A DNA which encodes the derivative of a monoclonal antibody or the derivative of the antibody fragment thereof which specifically reacts with ganglioside GD3 according to any one of (1) to (38).

(40) A recombinant vector comprising the DNA according to (39).

(41) A transformant which is obtained by introducing the recombinant vector according to (40) into a host cell.

(0031)

(42) A transformant KM871hIL2 (FERM BP-6918) which produces the antibody according to (35).

(43) A transformant KM8871hIL2 (FERM BP-6791) which produces the antibody according to (37).

(44) A process for producing an antibody, which comprises: culturing the transformant according to any one of (41) to (43) in a culture medium to produce and accumulate the derivative of a monoclonal antibody or the derivative of the antibody fragment thereof according to any one of (1) to (38) in the culture; and recovering the derivative of the antibody or the derivative of the antibody fragment thereof from the culture.

(0032)

(45) A human CDR-grafted antibody or the antibody fragment thereof which specifically reacts with ganglioside GD3.

(46) The human CDR-grafted antibody or the antibody fragment thereof according to (45), wherein the human CDR-grafted antibody comprises CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3.

(0033)

(47) The human CDR-grafted antibody or the antibody fragment thereof according to (45), wherein the human CDR-grafted antibody comprises: CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3; and FRs of an H chain V region and an L chain V region of a human antibody.

(48) The human CDR-grafted antibody or the antibody fragment thereof according to (45), wherein the human CDR-grafted antibody comprises: CDRs of an H chain V region and an L chain V region of a monoclonal antibody against ganglioside GD3; FRs of an H chain V region and an L chain V region of a human antibody; and an H chain C region and an L chain C region of a human antibody.

(0034)

(49) The human CDR-grafted antibody or the antibody fragment thereof according to any one of (46) to (48), wherein the antibody comprises CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively.

(50) The human CDR-grafted antibody or the antibody fragment thereof according to any one of (46) to (48), wherein the antibody comprises CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8, respectively.

(0035)

(51) The human CDR-grafted antibody or the antibody fragment thereof according to any one of (46) to (48), wherein the antibody comprises: CDR1, CDR2 and CDR3 of the H chain V region having the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, respectively; and CDR1, CDR2 and CDR3 of the L chain V region having the amino acid sequences represented by SEQ ID NOs:6, 7 and 8.

(52) The human CDR-grafted antibody or the antibody fragment thereof according to any one of (46) to (48), wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9.

(0036)

(53) The human CDR-grafted antibody or the antibody fragment thereof according to any one of (46) to (48), wherein the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

(54) The human CDR-grafted antibody or the antibody fragment thereof according to any one of (46) to (48), wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and the L chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

(0037)

(55) The human CDR-grafted antibody KM8871 or the antibody fragment thereof according to any one of (46) to (48), wherein the H chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:9; and the L

chain V region of the antibody comprises the amino acid sequence represented by SEQ ID NO:54.

(56) A DNA which encodes the human CDR-grafted antibody or the antibody fragment thereof which specifically reacts with ganglioside GD3 according to any one of (46) to (48).

(0038)

(57) A recombinant vector comprising the DNA according to (56).

(58) A transformant which is obtained by introducing the recombinant vector according to (57) into a host cell.

(59) A transformant KM8871 (FERM BP-6790) which produces the human CDR-grafted antibody according to (55).

(0039)

(60) A process for producing an antibody, which comprises: culturing the transformant according to (58) or (59) in a culture medium to produce and accumulate the human CDR-grafted antibody or the antibody fragment thereof according to any one of (45) to (55) in the culture; and recovering the antibody or the antibody fragment thereof from the culture.

(61) A therapeutic agent for cancers, comprising, as an active ingredient, at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to any one of (1) to (38) and the human CDR-grafted antibody and the antibody fragment thereof according to any one of (45) to (55).

(0040)

(62) A diagnostic agent for cancers, comprising, as an active ingredient, at least one selected from the derivative of a monoclonal antibody and the derivative of the antibody fragment thereof according to any one of (1) to (38) and the human CDR-grafted antibody and the antibody fragment thereof according to any one of (45) to (55).

(0041)

(Embodiments for carrying out the invention)

The present invention relates to derivatives of an antibody, comprising a monoclonal antibody which specifically reacts with ganglioside GD3 or the antibody fragment thereof which is conjugated with a radioisotope, a protein or a low molecular weight agent.

Examples of the monoclonal antibody include an antibody produced by a hybridoma, a humanized antibody, a human antibody and the like.

(0042)

A hybridoma is a cell, which is obtained by subjecting a B cell prepared by immunizing a non-human mammal with an antigen to cell fusion with a myeloma cell derived from a mouse or the like which produces a desired monoclonal antibody having an antigen specificity and.

5 Examples of the humanized antibody include a human chimeric antibody, a human CDR-grafted antibody and the like.
(0043)

10 A human chimeric antibody is an antibody which comprises an H chain V region (hereinafter referred to as "HV" or "VH") and an L chain V region (hereinafter referred to as "LV" or "VL") of an antibody derived from a non-human animal, an H chain C region (hereinafter referred to as "CH") of a human antibody and an L chain C region (hereinafter referred to as "CL") of a human antibody. As the non-human animal, any animal such as mouse, rat, hamster, rabbit or the like can be used, so long as a hybridoma can be prepared therefrom.

15 (0044)

20 The human chimeric antibody of the present invention can be produced by obtaining cDNAs encoding VH and VL from a hybridoma which produces a monoclonal antibody which specifically reacts with GD3, constructing a human chimeric antibody expression vector by inserting them into an expression vector for animal cell having genes encoding human antibody CH and human antibody CL, and expressing the expression vector by introducing it into an animal cell.

(0045)

25 As the CH of a human chimeric antibody, it may be any region which belongs to human immunoglobulin (hereinafter referred to as "hIg"), but those of hIgG class are suitable and any one of subclasses belonging to hIgG class, such as hIgG1, hIgG2, hIgG3 and hIgG4, can also be used. Also, as the CL of a human chimeric antibody, it may be any region which belongs to hIg, and those of κ class or λ class can be used.

(0046)

30 Examples of the anti-GD3 chimeric antibody include KM871 described in Japanese Published Unexamined Patent Application No. 304989/93.

A human CDR-grafted antibody is an antibody in which amino acid sequences of CDRs in VH and VL derived from a non-human animal antibody are grafted to appropriate positions of VH and VL of a human antibody.

35 The anti-GD3 CDR-grafted antibody of the present invention can be produced by constructing cDNAs encoding V regions in which CDR sequences of VH

and VL of an antibody derived from a non-human animal which specifically reacts with GD3 are grafted to CDR sequences of VH and VL of an optional human antibody, constructing a human CDR-grafted antibody expression vector by inserting them into an expression vector for animal cell having genes encoding human antibody CH and human antibody CL, and expressing the human CDR-grafted antibody by introducing the expression vector into an animal cell.

(0047)

As the CH of the human CDR-grafted antibody of the present invention, it may be any region which belongs to hIg, but those of hIgG class are suitable and any one of subclasses belonging to hIgG class, such as hIgG1, hIgG2, hIgG3 and hIgG4, can also be used. Also, as the CL of the human CDR-grafted antibody, it may be any region which belongs to hIg, and those of κ class or λ class can be used.

(0048)

Examples of the anti-GD3 CDR-grafted antibody include antibody KM8871 in which VH of the antibody comprises the amino acid sequence represented by SEQ ID NO:9 and CH thereof comprises an amino acid sequence of a human antibody IgG1 subclass, and VL of the antibody comprises the amino acid sequence represented by SEQ ID NO:54 and CL thereof comprises an amino acid sequence of a human antibody κ class.

Originally, a human antibody is an antibody naturally present in the human body. It also includes antibodies obtained from a human antibody phage library and a human antibody-producing transgenic animal which are produced based on the recent advance in genetic engineering, cell engineering and developmental engineering techniques.

(0049)

With regard to the antibody present in the human body, a lymphocyte capable of producing the antibody can be cultured by isolating a human peripheral blood lymphocyte and infecting it with EB virus or the like to thereby immortalizing it, followed by cloning, and the antibody can be purified from the culture mixture.

The human antibody phage library is a library in which antibody fragments such as Fab, scFv and the like are expressed on the phage surface by inserting an antibody gene prepared from human B cell into a phage gene. From the library, a phage expressing a desired antibody fragment having an antigen binding activity can be recovered, using its activity to bind to an antigen-immobilized substrate as the marker.

The antibody fragment can be genetic engineeringly converted further into a human antibody molecule having two complete H chains and two complete L chains.

(0050)

A human antibody-producing transgenic non-human animal is an animal in which a human antibody gene is introduced into cells. Specifically, a human antibody-producing transgenic animal can be produced by introducing a human antibody gene into mouse ES cell, transplanting the ES cell into an early stage embryo of other mouse and then developing it. As the preparation method of a human antibody from the human antibody-producing transgenic animal, the human antibody can be produced and accumulated in a culture mixture by obtaining a human antibody-producing hybridoma by a hybridoma production method usually carried out in non-human mammals and then culturing it.

(0051)

Examples of the antibody fragment include Fab, Fab', F(ab')₂, scFv, dsFv, a peptide comprising CDR and the like.

An Fab is an antibody fragment having a molecular weight of about 50,000 and antigen binding activity, in which about a half of the N-terminal side of H chain and the entire L chain, among fragments obtained by treating IgG with a protease, papain (cut at the 224th position amino acid residue of the H chain), are bound together through a disulfide bond.

(0052)

The Fab of the present invention can be obtained treating an antibody which specifically reacts with GD3 with a protease, papain. Also, the Fab can be produced by inserting DNA encoding Fab of the antibody into an expression vector for procaryote or an expression vector for eucaryote, and introducing the vector into a procaryote or eucaryote to express the Fab.

An F(ab')₂ is an antibody fragment having a molecular weight of about 100,000 and antigen binding activity, which is slightly larger than the Fab bound via a disulfide bond of the hinge region, among fragments obtained by treating IgG with a protease, pepsin (cut at the 234th position amino acid residue of the H chain).

(0053)

The F(ab')₂ of the present invention can be obtained treating an antibody which specifically reacts with GD3 with a protease, pepsin. Also, the F(ab')₂ can be produced by binding Fab' described below via a thioether bond or a disulfide bond.

An Fab' is an antibody fragment having a molecular weight of about 50,000 and antigen binding activity, which is obtained by cutting a disulfide bond of the hinge region of the F(ab')₂.

(0054)

The Fab' of the present invention can be obtained treating F(ab')₂ which specifically reacts with GD3 with a reducing agent, dithiothreitol. Also, the Fab' can be produced by inserting DNA encoding Fab' fragment of the antibody into an expression vector for procaryote or an expression vector for eucaryote, and introducing the vector into a procaryote or eucaryote to effect its expression.

(0055)

An scFv is a VH-P-VL or VL-P-VH polypeptide in which one chain VH and one chain VL are linked using an appropriate peptide linker (hereinafter referred to as "P"). The VH and VL in the scFv used in the present invention may be any antibody of an antibody produced by a hybridoma, a humanized antibody and a human antibody.

The scFv of the present invention can be produced by obtaining cDNA encoding the VH and VL of an antibody which specifically reacts with GD3, constructing DNA encoding scFv, inserting the DNA into an expression vector for procaryote or an expression vector for eucaryote, and then introducing the expression vector into a prokaryote or eucaryote to express the scFv.

(0056)

A dsFv is obtained by binding polypeptides in which one amino acid residue of each of VH and VL is substituted with a cysteine residue via a disulfide bond between the cysteine residues. The amino acid residue to be substituted with a cysteine residue can be selected based on a three-dimensional structure estimation of the antibody in accordance with the method shown by Reiter *et al.* (*Protein Engineering*, 7: 697 (1994)). The VH and VL in the dsFv of the present invention may be any antibody of an antibody produced by a hybridoma, a humanized antibody and a human antibody.

(0057)

The dsFv of the present invention can be produced by obtaining cDNA encoding the VH and VL of an antibody capable of specifically reacting with GD3, constructing DNA encoding dsFv, inserting the DNA into an expression vector for procaryote or an expression vector for eucaryote, and then introducing the expression vector into a prokaryote or eucaryote to express the dsFv.

A peptide comprising CDR is constituted by at least one region of H chain and L chain CDRs. Plural CDRs can be bound directly or via an appropriate peptide linker.

(0058)

The peptide comprising CDR of the present invention can be produced by obtaining cDNA encoding the VH and VL of an antibody capable of specifically reacting with GD3, inserting the cDNA into an expression vector for prokaryote or an expression vector for eucaryote, and then introducing the expression vector into a prokaryote or eucaryote to express the peptide comprising CDR.

The peptide comprising CDR can also be produced by a chemical synthesis method such as Fmoc method (fluorenylmethoxycarbonyl method) or tBoc method (t-butyloxycarbonyl method).

(0059)

The antibody derivatives of the present invention can be produced by chemically conjugating a radioisotope, a protein, a low molecular weight agent or the like to the N-terminal side or C-terminal side of the H chain or L chain of the antibody or the antibody fragment which specifically reacts with GD3, to an appropriate substituent or side chain of the antibody or the antibody fragment, or to a sugar chain of the antibody or the antibody fragment (*Antibody Engineering Handbook*, edited by Osamu Kanemitsu, published by Chijin Shokan (1994)).

(0060)

Also, the antibody derivatives can be genetically produced by linking a DNA encoding an antibody or the antibody fragment which specifically reacts with GD3 to other DNA encoding a protein to be bound, inserting the DNA into an expression vector, and introducing the expression vector into a host cell.

Examples of the isotope include ^{131}I , ^{125}I and the like, and they can be conjugated to an antibody by, e.g., a chloramine T method.

(0061)

Examples of the low molecular weight agent include anticancer agents such as alkylating agent (e.g., nitrogen mustard, cyclophosphamide, *etc.*), metabolic antagonist (e.g., 5-fluorouracil, methotrexate, *etc.*), antibiotic (e.g., daunomycin, bleomycin, mitomycin C, daunorubicin, doxorubicin, *etc.*), plant alkaloid (e.g., vincristine, vinblastine, vindesine, *etc.*), hormone drug (e.g., tamoxifen, dexamethasone, *etc.*) and the like (*Clinical Oncology*, edited by Japanese Society of Clinical Oncology, published by Cancer and Chemotherapy (1996)); anti-inflammatory agents such as steroid agent (e.g., hydrocortisone, prednisone, *etc.*), non-steroidal drug (e.g., aspirin, indometacin, *etc.*), immunomodulator (e.g., aurothiomalate, penicillamine, *etc.*), immunosuppressing agent (e.g., cyclophosphamide, azathioprine, *etc.*), antihistaminic agent (e.g., chlorpheniramine maleate, clemastine, *etc.*) and the like (*Inflammation and Anti-inflammatory Therapy*, Ishiyaku Shuppan (1982)); and the like. Examples of the

method for conjugating daunomycin to an antibody include a method in which daunomycin and an amino group of an antibody are conjugated via glutaraldehyde, a method in which an amino group of daunomycin and a carboxyl group of an antibody are conjugated via a water-soluble carbodiimide, and the like.

5 (0062)

As the protein, cytokine which activates immunocompetent cells is preferred, and examples include hIL-2, hGM-CSF, human macrophage colony-stimulating factor (hereinafter referred to as "hM-CSF"), human interleukin 12 (hereinafter referred to as "hIL-12") and the like. Also, in order to injure cancer cells directly, toxins such as
10 ricin, diphtheria toxin and the like can be used. For example, a fusion antibody with a protein can be produced by conjugating a cDNA encoding an antibody or the antibody fragment to other cDNA encoding the protein, constructing a DNA encoding the fusion antibody, inserting the DNA into an expression vector for prokaryote or an expression vector for eucaryote, and then introducing the expression vector into a prokaryote or
15 eucaryote to express the fusion antibody.

(0063)

Examples of the fusion protein of the anti-GD3 human antibody with cytokine include a fusion protein of anti-GD3 CDR-grafted antibody KM8871 with hIL-2 and a fusion protein of anti-GD3 chimeric antibody KM871 with hIL-2. Examples
20 of the fusion protein of anti-GD3 CDR-grafted antibody KM8871 with hIL-2 include KM8871-hIL-2 in which the H chain V region of the antibody conjugated with hIL-2 has the amino acid sequence represented by SEQ ID NO:53, the L chain V region of the antibody has the amino acid sequence represented by SEQ ID NO:54 and the L chain C region has an amino acid sequence of a human antibody κ class. Examples of the
25 fusion protein of anti-GD3 chimeric antibody KM871 with hIL-2 include KM871-hIL-2 in which the H chain V region of the antibody conjugated with hIL-2 has the amino acid sequence represented by SEQ ID NO:57, the L chain V region of the antibody has the amino acid sequence represented by SEQ ID NO:56 and the L chain C region has an amino acid sequence of a human antibody κ class.

30 The methods for producing a humanized antibody which specifically reacts with GD3 and derivatives of an antibody or the antibody fragment which specifically reacts with GD3.

(0064)

1. Production of humanized antibody

35 (1) Construction of humanized antibody expression vector

The humanized antibody expression vector is an expression vector for animal cell to which genes encoding the CH and CL of a human antibody are inserted, and can be constructed by cloning the genes which encode the CH and CL of a human antibody into an expression vector for animal cell. The C regions of a human antibody can be any human antibody CH and CL, and examples include a C region belonging to IgG1 subclass of in an H chain of a human antibody (hereinafter referred to as "hC γ 1") and a C region belonging to κ class in an L chain of a human antibody (hereinafter referred to as "hC κ "). A chromosomal DNA comprising an exon and an intron can be used as the gene encoding CH and CL of a human antibody, and cDNA can also be used. Any expression vector for animal cell can be used, so long as a gene encoding the human antibody C region can be inserted and expressed. Examples include pAGE107 (*Cytotechnology*, 3, 133 (1990)), pAGE103 (*J. Biochem.*, 101, 1307 (1987)), pHSG274 (*Gene*, 27, 223 (1984)), pKCR (*Proc. Natl. Acad. Sci. U.S.A.*, 78, 1527 (1981)), pSG1 β d2-4 (*Cytotechnology*, 4, 173 (1990)) and the like. Examples of the promoter and the enhancer used in the expression vector for animal cell include early promoter and enhancer of SV40 (*J. Biochem.*, 101, 1307 (1987)), LTR promoter and enhancer of Moloney mouse leukemia virus (*Biochem. Biophys. Res. Commun.*, 149, 960 (1987)), promoter (*Cell*, 41, 479, 1985)) and enhancer (*Cell*, 33, 717 (1983)) of immunoglobulin H chain and the like.

The humanized antibody expression vector may be either of a type in which a gene encoding an antibody H chain and a gene encoding an antibody L chain exist on separate vectors or of a type in which both genes exist on the same vector (tandem type). In respect of easiness of construction of a humanized antibody expression vector, easiness of introduction into animal cells, and balance between the expression levels of antibody H and L chains in animal cells, a tandem type of the humanized antibody expression vector is more preferable (*J. Immunol. Methods*, 167, 271 (1994)). Examples of the tandem type humanized antibody expression vector include pKANTEX93 (WO 97/10354), pEE18 (*HYBRIDOMA*, 17, 559 (1998)) and the like.

The thus constructed humanized antibody expression vector can be used for the expression of the human chimeric antibody and human CDR-grafted antibody in animal cells.

(0066)

(2) Preparation of cDNA encoding V region of antibody derived from non-human animal

cDNAs encoding VH and VL of an antibody derived from a non-human animal such as a mouse antibody are obtained as follows.

(0067)

mRNA is extracted from hybridoma cells producing a mouse antibody or the like to synthesize cDNA. The synthesized cDNA is inserted into a vector such as a phage, a plasmid or the like, to prepare a cDNA library. Each of a recombinant phage or recombinant plasmid containing cDNA encoding VH and a recombinant phage or recombinant plasmid containing cDNA encoding VL is isolated from the library using a C region part or a V region part of a mouse antibody as the DNA probe. The full nucleotide sequences of the VH and VL of the mouse antibody of interest on the recombinant phage or recombinant plasmid are determined, and the full amino acid sequences of the VH and VL are deduced from the nucleotide sequences.

(0068)

The non-human animal may be any animal such as mouse, rat, hamster, rabbit or the like, so long as a hybridoma cell can be produced therefrom.

Examples of the method for preparing total RNA from a hybridoma cell include a guanidine thiocyanate-cesium trifluoroacetate method (*Methods in Enzymol.*, 154, 3 (1987)) and the like. Examples of the method for preparing mRNA from total RNA include an oligo (dT) immobilized cellulose column method (*Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Lab. Press, New York (1989), hereinafter referred to as "*Molecular Cloning: A Laboratory Manual*") and the like. Also, examples of a kit for preparing mRNA from a hybridoma cell include Fast Track mRNA Isolation Kit (manufactured by Invitrogen), Quick Prep mRNA Purification Kit (manufactured by Pharmacia) and the like.

(0069)

Examples of the method for synthesizing cDNA and preparing a cDNA library include known methods (*Molecular Cloning: A Laboratory Manual*; *Current Protocols in Molecular Biology*, Supplement 1-34); a method using a commercially available kit such as Super Script™ Plasmid System for cDNA Synthesis and Plasmid Cloning (manufactured by GIBCO BRL), ZAP-cDNA Kit (manufactured by Stratagene), etc.; and the like.

(0070)

The vector into which the cDNA synthesized using mRNA extracted from a hybridoma cell as the template is inserted for preparing a cDNA library may be any vector, so long as the cDNA can be inserted. Examples include ZAP Express (*Strategies*, 5: 58 (1992)), pBluescript II SK(+) (*Nucleic Acids Research*, 17: 9494

(1989)), λ zapII (manufactured by Stratagene), λ gt10 and λ gt11 (*DNA Cloning: A Practical Approach*, I, 49 (1985)), Lambda BlueMid (manufactured by Clontech), λ ExCell and pT7T3 18U (manufactured by Pharmacia), pcD2 (*Mol. Cell. Biol.*, 3: 280 (1983)), pUC18 (*Gene*, 33: 103 (1985)), and the like.

5 (0071)

Any *E. coli* for introducing the cDNA library constructed by a phage or plasmid vector may be used, so long as the cDNA library can be introduced, expressed and maintained. Examples include XL1-Blue MRF' (*Strategies*, 5: 81 (1992)), C600 (*Genetics*, 39: 440 (1954)), Y1088 and Y1090 (*Science*, 222: 778 (1983)), NMS22 (*J. Mol. Biol.*, 166: 1 (1983)), K802 (*J. Mol. Biol.*, 16: 118 (1966)), JM105 (*Gene*, 38: 275 (1985)), and the like.

(0072)

For selecting cDNA clones encoding VH and VL of an antibody derived from a non-human animal in the cDNA library, a colony hybridization or plaque hybridization method using an isotope- or fluorescence-labeled probe may be used (*Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Lab. Press, New York (1989)). Also, the cDNAs encoding VH and VL can be prepared through polymerase chain reaction (hereinafter referred to as "PCR"; *Molecular Cloning: A Laboratory Manual*; *Current Protocols in Molecular Biology*) by preparing primers and using cDNA prepared from mRNA or a cDNA library as the template.

(0073)

The nucleotide sequence of the cDNA can be determined by digesting the cDNA selected by the above method with appropriate restriction enzymes and the like, cloning the fragments into a plasmid such as pBluescript SK(-) (manufactured by Stratagene) or the like, carrying out the reaction by a usually used nucleotide analyzing method such as the dideoxy method of Sanger, F. *et al.* (*Proc. Natl. Acad. Sci. USA*, 74: 5463 (1977)) or the like, and then analyzing the sequence using an automatic nucleotide sequence analyzer such as A.L.F. DNA sequencer (manufactured by Pharmacia) or the like.

(0074)

Whether the obtained cDNAs encode the full amino acid sequences of the VH and VL of the antibody containing a secretory signal sequence can be confirmed by estimating the full amino acid sequences of the VH and VL from the determined nucleotide sequence and comparing them with full amino acid sequences of the VH and VL of known antibodies (*Sequences of Proteins of Immunological Interest*, US Dept.

Health and Human Services (1991), hereinafter referred to as "*Sequences of Proteins of Immunological Interest*").

(0075)

(3) Analysis of amino acid sequence of V region derived from a non-human animal

5 Regarding the complete amino acid sequences of VH and VL of the antibody comprising a secretory signal sequence, the length of the secretory signal sequence and N-terminal amino acid sequences can be deduced and subgroups to which they belong can also be known, by comparing with full amino acid sequences of VH and VL of known antibodies (*Sequences of Proteins of Immunological Interest*). Also, regarding the amino acid sequence of each CDR of VH and VL, it can be found by comparing it with amino acid sequences of VH and VL of known antibodies (*Sequences of Proteins of Immunological Interest*).

(0076)

(4) Construction of human chimeric antibody expression vector

15 A human chimeric antibody expression vector can be constructed by cloning cDNAs encoding VH and VL of an antibody derived from a non-human animal in the region upstream of genes encoding CH and CL of the human antibody on the humanized antibody expression vector as described in the item 1(1). For example, each of cDNAs encoding VH and VL of an antibody derived from a non-human animal is linked with a synthesized DNA comprising nucleotide sequences at the 3' terminals of VH and VL of an antibody derived from a non-human animal and nucleotide sequences at the 5' terminals of CH and CL of a human antibody and having a recognition sequence of an appropriate restriction enzyme at both terminals, and each cDNA is cloned so that it is appropriately expressed in upstream of genes encoding the CH and CL of the humanized antibody expression vector as described in the item 1(1) to thereby construct a human chimeric antibody expression vector.

(0077)

(5) Construction of cDNA encoding V region of human CDR-grafted antibody

30 cDNAs encoding VH and VL of a human CDR-grafted antibody can be obtained as follows. First, amino acid sequences of FRs in VH and VL of a human antibody to which amino acid sequences of CDRs in VH and VL of an antibody derived from a non-human animal antibody are grafted are selected. Any amino acid sequences of FRs in VH and VL of a human antibody can be used, so long as they are derived from human. Examples include amino acid sequences of FRs in VH and VL of human antibodies registered in database such as Protein Data Bank or the like, and amino acid sequences common to subgroups of FRs in the VH and VL of human

antibodies (*Sequences of Proteins of Immunological Interest*) and the like. In order to produce a human CDR-grafted antibody having potent activity, amino acid sequences having high homology (at least 60% or more) with amino acid sequence of FRs of VH and VL of an antibody of interest derived from a non-human animal is preferably selected. Then, amino acid sequences of CDRs of VH and VL of the antibody derived from a non-human animal are grafted to the selected amino acid sequences of FRs of VH and VL of a human antibody to design amino acid sequences of the VH and VL of a human CDR-grafted antibody. The designed amino acid sequences are converted to DNA sequences by considering the frequency of codon usage found in nucleotide sequences of genes of antibodies (*Sequence of Proteins of Immunological Interest*), and the DNA sequences encoding the amino acid sequences of the VH and VL of a human CDR-grafted antibody are designed. Several synthetic DNAs having a length of about 100 nucleotides are synthesized, and PCR is carried out using them. In this case, it is preferred in each of the H chain and the L chain that 6 synthetic DNAs are designed in view of the reaction efficiency of PCR and the lengths of DNAs which can be synthesized.

(0078)

Furthermore, they can be easily cloned into the humanized antibody expression vector constructed in the item 1(1) by introducing the recognition sequence of an appropriate restriction enzyme to the 5' terminal of the synthetic DNAs present on the both terminals. After the PCR, an amplified product is cloned into a plasmid such as pBluescript SK (-) (manufactured by Stratagene) or the like, and the nucleotide sequences are determined according to the method described in the item 1(2) to obtain a plasmid having DNA sequences encoding the VH and VL of a designed human CDR-grafted antibody.

(0079)

(6) Modification of amino acid sequence of V region of human CDR-grafted antibody

It is known that when a human CDR-grafted antibody is produced by simply grafting only CDRs in VH and VL an antibody derived from a non-human animal in FRs of the VH and VL of a human antibody, its antigen binding activity is reduced in comparison with that of the original antibody derived from a non-human animal (*BIO/TECHNOLOGY*, 9, 266 (1991)). As the reason, it is considered that several amino acid residues in not only CDRs but also FRs directly or indirectly relate to the antigen binding activity in the VH and VL of the original antibody derived from a non-human animal, and that these amino acid residues are changed to different amino acid residues derived from FRs of the VH and VL of the human antibody. In order to

resolve the problem, in human CDR-grafted antibodies, attempts have been made to identify, among amino acid sequences of the FR of the VH and VL of human antibodies, an amino acid residue which directly relates to binding to the antibody, an amino acid residue which is interacted with an amino acid residue of CDR or an amino acid residue which keeps three-dimensional structure of the antibody and is directly related to its binding to the antigen, and to increase the reduced antigen binding activity by changing them into amino acid residues of the original antibody derived from a non-human animal (*BIO/TECHNOLOGY*, 9, 266 (1991)). In producing a human CDR-grafted antibody, it is the most important point to efficiently identify these FR amino acid residues related to the antigen binding activity, so that construction and analysis of the three-dimensional structure of antibodies have been carried out by X-ray crystallography (*J. Mol. Biol.*, 112, 535 (1977)), computer-modeling (*Protein Engineering*, 7, 1501 (1994)) and the like. Although the information of the three-dimensional structure of antibodies has been useful in the production of a human CDR-grafted antibody, no method for producing a human CDR-grafted antibody which can be applied to any antibodies has been established yet. Therefore, various attempts must be currently be necessary, for example, several modified antibodies of each antibody are produced and the relationship between each of the modified antibodies and its antibody binding activity is examined.

(0080)

Substitution of the amino acid residues of the FR in the VH and VL of a human antibody can be achieved by carrying out the PCR described in the item 1(5) using synthetic DNA for modification. With regard to an amplified product obtained by the PCR, the nucleotide sequence is determined according to the method described in the item 1(2) so that whether the objective modification has been carried out is confirmed.

(0081)

(7) Construction of human CDR-grafted antibody expression vector

A human CDR-grafted antibody expression vector can be constructed by cloning cDNAs encoding VH and VL of the human CDR-grafted antibody constructed in the items 1(5) and 1(6) into upstream of the genes encoding CH and CL of the human antibody in the humanized antibody expression vector as described in the item 1(1).

(0082)

For example, when recognition sites for an appropriate restriction enzymes are introduced to the 5'-terminal of synthetic DNAs positioned at both terminals among synthetic DNAs used in the construction of VH and VL of the human CDR-grafted

antibody in the items 1(5) and 1(6), cloning can be carried out so that they are expressed in an appropriate form in upstream of genes encoding CH and CL of the human antibody in the humanized antibody expression vector as described in the item 1(1).

(0083)

5 (8) Transient expression of humanized antibody

In order to efficiently evaluate the antigen binding activity of various humanized antibodies produced, the humanized antibodies can be expressed transiently using the humanized antibody expression vector as described in the items 1(4) and 1(7) or the modified expression vector thereof. Any cell can be used as a host cell, so long as the host cell can express a humanized antibody. Generally, COS-7 cell (ATCC CRL1651) is used in view of its high expression level (*Methods in Nucleic Acids Res.*, CRC Press, 283 (1991)). Examples of the method for introducing the expression vector into COS-7 cell include a DEAE-dextran method (*Methods in Nucleic Acids Res.*, CRC Press, 283 (1991)), a lipofection method (*Proc. Natl. Acad. Sci. USA*, 84, 7413 (1987)), and the like.

(0084)

After introduction of the expression vector, the expression level and antigen binding activity of the humanized antibody in the culture supernatant can be measured by enzyme immunoassay (hereinafter referred to as "ELISA"; *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Chapter 14 (1988), hereinafter referred to as "*Antibodies: A Laboratory Manual*", *Monoclonal Antibodies: Principles and Practice*, Academic Press Limited (1996), hereinafter referred to as "*Monoclonal Antibodies*" (1996)) and the like.

(0085)

25 (9) Stable expression of humanized antibody

A transformant which produces a humanized antibody stably can be obtained by introducing into an appropriate host cell the humanized antibody expression vector described in the items 1(4) and 1(7).

Examples of the method for introducing the expression vector into a host cell include electroporation (Japanese Published Unexamined Patent Application No. 257891/90, *Cytotechnology*, 3, 133 (1990)) and the like.

(0086)

Any cell can be used as the host cell into which the humanized antibody expression vector is to be introduced, so long as it can express a humanized antibody. Examples include mouse SP2/0-Ag14 cell (ATCC CRL1581), mouse P3X63-Ag8.653 cell (ATCC CRL1580), CHO cell in which a dihydrofolate reductase gene (hereinafter

referred to as "DHFR gene") is detective (*Proc. Natl. Acad. Sci. U.S.A.*, 77: 4216 (1980)), rat YB2/3HL.P2.G11.16Ag.20 cell (ATCC CRL1662, hereinafter referred to as "YB2/0 cell"), and the like. The YB2/0 cell is preferable, since ADCC activity of the humanized antibody is enhanced when expressed therein.

5 (0087)

After introduction of the expression vector, transformants which express a humanized antibody stably are selected in accordance with the method disclosed in Japanese Published Unexamined Patent Application No. 257891/90, by culturing in a medium for animal cell culture containing an agent such as G418 sulfate (hereinafter referred to as "G418", manufactured by Sigma) or the like. Examples of the medium for animal cell culture include PRMI1640 medium (manufactured by Nissui Pharmaceutical), GIT medium (manufactured by Nissui Pharmaceutical), EX-CELL302 medium (manufactured by JRH), IMDM medium (manufactured by GIBCO BRL), Hybridoma-SFM medium (manufactured by GIBCO BRL), media obtained by adding various additives such as fetal bovine serum (hereinafter referred to as "FBS") to these media, and the like. The humanized antibody can be produced and accumulated in a culture medium by culturing the selected transformants in a medium. The expression level and antigen binding activity of the humanized antibody in the culture supernatant can be measured by ELISA or the like. Also, in the transformant, the expression level of the humanized antibody can be increased by using dhfr amplification system or the like according to the method disclosed in Japanese Published Unexamined Patent Application No. 257891/90.

20 (0088)

The humanized antibody can be purified from the culture supernatant of the transformant by using a protein A column (*Antibodies, A Laboratory Manual, Monoclonal Antibodies* (1996)). Any other conventional methods for protein purification can be used. For example, the humanized antibody can be purified by a combination of gel filtration, ion-exchange chromatography, ultrafiltration and the like. The molecular weight of the H chain or the L chain of the purified humanized antibody or the antibody molecule as a whole is determined by polyacrylamide gel electrophoresis (hereinafter referred to as "SDS-PAGE") (*Nature*, 227: 680 (1970)), Western blotting (*Antibodies, A Laboratory Manual, Monoclonal Antibodies* (1996)), and the like.

30 (0089)

35 (10) Evaluation of activity of humanized antibody

The binding activity to an antigen and the binding activity to cultured cancer cell lines of the purified humanized antibody can be measured by ELISA, an immunofluorescent method (*Cancer Immunol. Immunother.*, 36, 373 (1993)) and the like means. The cytotoxic activity against an antigen positive culture cell line can be evaluated by measuring the CDC activity, the ADCC activity or the like (*Cancer Immunol. Immunother.*, 36, 373 (1993)).

(0090)

(11) Application method of humanized antibody

It is considered to be useful in the diagnosis and treatment of human cancers and the like, such as lung cancer, melanoma, glioma, neuroblastoma, *etc.*, since the humanized antibody specifically binds to GD3 which expresses in cultured cancer cell lines derived from human and shows cytotoxic activities such as CDC activity, ADCC activity and the like. In addition, since the majority of the antibody are derived from the amino acid sequence of a human antibody in comparison with antibodies of non-human animals, it is expected that it will show strong anti-tumor effects in the human body without showing immunogenicity, and the therapeutic effects will be maintained for a prolonged period of time.

(0091)

The humanized antibody of the present invention can be administered alone, but it is generally preferred to provide it in the form of a pharmaceutical formulation produced by mixing it with at least one pharmaceutically acceptable carrier in accordance with a method well known in the technical field of pharmaceuticals.

It is preferred to select a route of administration which is the most effective in carrying out the intended treatment such as oral administration or parenteral administration, e.g., intraoral administration, tracheal administration, rectal administration, subcutaneous injection, intramuscular injection, intravenous injection, and the like. Intravenous injection is preferred in an antibody formulation.

(0092)

The dosage form includes sprays, capsules, tablets, granules, syrups, emulsions, suppositories, injections, ointments, tapes, and the like.

Examples of formulation suitable for oral administration include emulsions, syrups, capsules, tablets, powders, granules, and the like.

Liquid preparations such as emulsions and syrups, can be produced using additives such as water, saccharides, e.g., sucrose, sorbitol, fructose, *etc.*, glycols, e.g., polyethylene glycol, propylene glycol, *etc.*, oils, e.g., sesame oil, olive oil, soybean oil,

etc., antiseptics, e.g., p-hydroxybenzoate, *etc.*, flavors, e.g., strawberry flavor, peppermint, *etc.*, and the like.

(0093)

Capsules, tablets, powders, granules and the like can be produced using additives such as fillers, e.g., lactose, glucose, sucrose, mannitol, *etc.*, disintegrating agents, e.g., starch, sodium alginate, *etc.*, lubricants, e.g., magnesium stearate, *etc.*, binders, e.g., polyvinyl alcohol, hydroxypropylcellulose, gelatin, *etc.*, surfactants, e.g., fatty acid ester, *etc.*, plasticizers, e.g., glycerine, *etc.*, and the like.

(0094)

Examples of formulations suitable for parenteral administration include injections, suppositories, sprays, and the like.

Injections can be prepared using a carrier such as a salt solution, glucose solution or a mixture thereof, or the like.

Suppositories can be prepared using a carrier such as cacao butter, hydrogenated fat, a carboxylic acid, or the like.

Also, sprays can be prepared from the antibody or peptide itself or using a carrier or the like which does not stimulate oral and airway mucous membranes of patients and can facilitate absorption of the antibody or peptide by dispersing it as minute particles.

(0095)

Examples of the carrier include lactose, glycerine, and the like. Depending on the properties of the antibody or peptide and the carrier to be used, aerosols, dry powders and the like can be produced. The additives exemplified in the oral preparations can also be added to the parenteral preparations.

The dose and frequency of administration vary depending on intended therapeutic effect, administration method, treating period, age, body weight and the like, but the dose is generally from 10 µg/kg to 8 mg/kg per day per adult.

(0096)

2. Production of fusion protein of antibody or antibody fragment with cytokine

(1) Construction of gene encoding fusion protein of antibody or antibody fragment with cytokine

A gene encoding a fusion protein of an antibody or the antibody fragment with cytokine can be constructed by linking a gene encoding cytokine to the 5'-terminal or 3'-terminal of a gene encoding the H chain or L chain of an antibody or the antibody fragment via an appropriate synthetic DNA. Also, a gene encoding a fusion protein of an antibody or the antibody fragment with cytokine can also be constructed by

introducing an appropriate restriction enzyme recognition sequence into the 5'-terminal of a primer for amplification when a gene encoding cytokine is amplified by PCR, and linking the resulting gene to the 5'-terminal or 3'-terminal of a gene encoding the H chain or L chain of an antibody or the antibody fragment. Any chromosomal DNA or cDNA can be used as the gene encoding cytokine. The nucleotide sequence of the thus constructed gene encoding a fusion protein of an antibody or antibody fragment with cytokine is determined by the method described in the item 1(2) to confirm that it is the sequence of interest.

(0097)

(2) Construction of expression vector of fusion protein of antibody or antibody fragment with cytokine

An expression vector of a fusion protein of an antibody or the antibody fragment with cytokine can be constructed by substituting a part or whole of a gene encoding the H chain or L chain of an antibody or the antibody fragment on the humanized antibody expression vector described in the items 1(4) and 1(7) by the gene encoding a fusion protein of an antibody or the antibody fragment with cytokine described in the item 2(1). For example, when a fusion protein in which cytokine is conjugated with the C-terminal of the H chain of an antibody is prepared, the expression vector can be constructed by linking a gene encoding cytokine to the 3'-terminal of a gene encoding the CH of an antibody or the antibody fragment to thereby construct a gene encoding a fusion protein of the CH of an antibody or the antibody fragment with cytokine according to the item 2(1), and substituting, with the gene, a gene encoding the CH of an antibody or the antibody fragment on the humanized antibody expression vector described in the items 1(4) and 1(7).

(0098)

(3) Stable expression of fusion protein of antibody or antibody fragment with cytokine

Stable expression of a fusion protein of an antibody or the antibody fragment with cytokine in accordance with the method described in the item 1(9) using the expression vector of a fusion protein of an antibody or the antibody fragment with cytokine described in the item 2(2) is carried out so that a transformant capable of stably expressing the fusion protein of an antibody or the antibody fragment with cytokine can be obtained, the fusion protein of an antibody or antibody fragment with cytokine is purified from its culture supernatant, and its molecular weight and the like can be analyzed.

(0099)

(4) Evaluation of *in vitro* activity of fusion protein of antibody or antibody fragment with cytokine

Among activities of the thus purified fusion protein of an antibody or the antibody fragment with cytokine, activities of the antibody moiety, namely binding activity with an antigen and its binding activity to cultured cancer cell lines can be measured by ELISA, an immunofluorescent method and the like. Also, its cytotoxic activities against antigen-positive cultured cancer cell lines can be evaluated by measuring its CDC activity, ADCC activity and the like. On the other hand, activity of the cytokine moiety can be evaluated, e.g., using growth-supporting activity of a cultured cell line which shows concentration-dependent growth against cytokine, as an index (*Proc. Natl. Acad. Sci. U.S.A.*, 91, 9626 (1994)).

(0100)

(5) Evaluation of *in vivo* activity of fusion protein of an antibody or antibody fragment with cytokine

Antitumor effect of the fusion protein of an antibody or the antibody fragment with cytokine can be evaluated by administering to a syngenic graft mouse model in which an antigen-expressing cultured mouse cancer cell line is transplanted into a wild type mouse having normal immune system. Also, its antitumor effect in the living body can be evaluated by comparing respective cases of the administration of the antibody alone, cytokine alone and simultaneous administration of the antibody and cytokine to the syngenic graft mouse model (*Proc. Natl. Acad. Sci. U.S.A.*, 93, 7826 (1996)). To date, there are no reports on GD2- or GD3-positive mouse culture cell lines, which are specifically expressed in human melanoma and neuroblastoma. However, a GD2- or GD3-positive transformant can be produced by introducing a ganglioside synthase gene of interest to be specifically expressed into a mouse culture cell line which expresses GM2 or GM3 as a biosynthesis precursor of GD2 or GD3 (*Proc. Natl. Acad. Sci. U.S.A.*, 91, 10455 (1994)). A syngenic graft model can be produced by transplanting this transformant into a wild type mouse having a normal immune system.

The thus prepared mouse model can be used in the *in vivo* evaluation of a fusion protein of an antibody or the antibody fragment with cytokine against a ganglioside of interest.

(0101)

(6) Application method of fusion proteins of humanized antibody or antibody fragment with cytokine

Since the fusion protein of an antibody or the antibody fragment with cytokine of the present invention specifically binds to GD3 expressing in cultured cancer cell lines of human origin and shows cytotoxic activities such as CDC activity, ADCC activity and the like, it is considered to be useful in the diagnosis and treatment of human cancers and the like such as lung cancer, melanoma, glioma, neuroblastoma, etc. Since the majority of the humanized antibody or antibody fragment of the present invention are derived from the amino acid sequence of a human antibody in comparison with antibodies of non-human animals, it is expected that it will show strong anti-tumor effects in the human body without showing immunogenicity, and the therapeutic effects will be maintained for a prolonged period of time, and since it can activate immunocompetent cells in the periphery of a cancer by the activity of the conjugated cytokine moiety, stronger antitumor effects are expected than the administration of the antibody alone, cytokine alone or simultaneous administration of the antibody and cytokine, and reduction of side effects is expected in comparison with the systemic administration of cytokine.

(0102)

The fusion protein of a humanized antibody or the antibody fragment with cytokine of the present invention can be administered alone, but it is generally preferred to provide it in the form of a pharmaceutical formulation produced by mixing it with at least one pharmaceutically acceptable carrier in accordance with a method well known in the technical field of pharmaceuticals.

It is preferred to select a route of administration which is the most effective in carrying out the intended treatment such as oral administration or parenteral administration, e.g., intraoral administration, tracheal administration, rectal administration, subcutaneous injection, intramuscular injection, intravenous injection, and the like. Intravenous injection is preferred in a protein formulation.

(0103)

The dosage form includes sprays, capsules, tablets, granules, syrups, emulsions, suppositories, injections, ointments, tapes, and the like.

Examples of formulation suitable for oral administration include emulsions, syrups, capsules, tablets, powders, granules, and the like.

Liquid preparations such as emulsions and syrups, can be produced using additives such as water, saccharides, e.g., sucrose, sorbitol, fructose, etc., glycols, e.g., polyethylene glycol, propylene glycol, etc., oils, e.g., sesame oil, olive oil, soybean oil, etc., antiseptics, e.g., p-hydroxybenzoate, etc., flavors, e.g., strawberry flavor, peppermint, etc., and the like.

(0104)

Capsules, tablets, powders, granules and the like can be produced using additives such as fillers, e.g., lactose, glucose, sucrose, mannitol, *etc.*, disintegrating agents, e.g., starch, sodium alginate, *etc.*, lubricants, e.g., magnesium stearate, *etc.*, binders, e.g., polyvinyl alcohol, hydroxypropylcellulose, gelatin, *etc.*, surfactants, e.g., fatty acid ester, *etc.*, plasticizers, e.g., glycerine, *etc.*, and the like.

(0105)

Examples of formulations suitable for parenteral administration include injections, suppositories, sprays, and the like.

Injections can be prepared using a carrier such as a salt solution, glucose solution or a mixture thereof, or the like.

Suppositories can be prepared using a carrier such as cacao butter, hydrogenated fat, a carboxylic acid, or the like.

Also, sprays can be prepared from the antibody or peptide itself or using a carrier or the like which does not stimulate oral and airway mucous membranes of patients and can facilitate absorption of the antibody or peptide by dispersing it as minute particles.

(0106)

Examples of the carrier include lactose, glycerine, and the like. Depending on the properties of the antibody or peptide and the carrier to be used, aerosols, dry powders and the like can be produced. The additives exemplified in the oral preparations can also be added to the parenteral preparations.

The dose and frequency of administration vary depending on intended therapeutic effect, administration method, treating period, age, body weight and the like, but the dose is generally from 10 µg/kg to 8 mg/kg per day per adult.

(0107)

Examples of the present invention are shown below.

(0108)

(Examples)

Example 1. Production of anti-GD3 CDR-grafted antibody:

1. Analysis of amino acid sequence of V region of anti-GD3 mouse antibody KM641

A full amino acid sequence of VH of anti-GD3 mouse antibody KM641 disclosed in Japanese Published Unexamined Patent Application No. 304989/93 is represented by SEQ ID NO:1, and a full amino acid sequence of VL thereof is represented by SEQ ID NO:2. Based on the comparison of both sequences with analytical results of amino acid sequences of known antibodies (*Sequences of Proteins*

of Immunological Interest) and N-terminal amino acid sequences of H chain and L chain of purified anti-GD3 mouse antibody KM641, it was confirmed that positions (-19) to (-1) in the amino acid sequence of H chain and positions (-20) to (-1) in the amino acid sequence of L chain are secretory signal sequences. Full amino acid sequences of the secretory VH and VL are represented by SEQ ID NO:55 and SEQ ID NO:56, respectively. Also, it was found that CDRs 1, 2 and 3 of VH have the amino acid sequences represented by SEQ ID NOs:3, 4 and 5, and CDRs 1, 2 and 3 of VL have those represented by SEQ ID NOs:6, 7 and 8.

(0109)

2. Measurement of binding activity of antibody to various ganglioside (ELISA)

The binding activity of antibodies to various gangliosides was measured as follows.

Each ganglioside (2 nmol) was dissolved in 2 ml of ethanol solution containing 10 µg of dipalmitoylphosphatidylcholine (manufactured by SIGMA) and 5 µg of cholesterol (manufactured by SIGMA). In each well of a 96 well plate for ELISA use (manufactured by Greiner), 20 µl of the solution (becomes 20 pmol/well) or 20 µl of the solution diluted with ethanol was respectively dispensed, followed by air-drying, and then PBS containing 1% BSA (hereinafter referred to as "1% BSA-PBS") was added in 100 µl/well and allowed to react at room temperature for 1 hour to thereby block remaining active groups. After discarding 1% BSA-PBS, culture supernatants of transformants or diluted solutions of humanized antibodies were added in 50 µl/well and allowed to react at room temperature for 1 hour. After the reaction, each well was washed with PBS containing 0.05% Tween 20 (hereinafter referred to as "Tween-PBS") and then a peroxidase-labeled goat anti-human IgG(γ) antibody solution (manufactured by Kirkegaard & Perry Laboratories) diluted 1,000 folds with 1% BSA-PBS was added as a secondary antibody solution in 50 µl/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, an ABTS substrate solution (a solution prepared by dissolving 0.55 g of 2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid) in 1 L of 0.1 M citrate buffer (pH 4.2) and adding 1 µl/ml hydrogen peroxide just before the use) was added in 50 µl/well for color development and the absorbance at 415 nm (hereinafter referred to as "OD415") was measured.

(0110)

3. Construction of cDNAs encoding VH and VL of anti-GD3 CDR-grafted antibody

(1) Design of amino acid sequences of VH and VL of anti-GD3 CDR-grafted antibody

First, an amino acid sequence of VH of an anti-GD3 CDR-grafted antibody was designed as follows. In order to graft the CDR amino acid sequences of VH of the anti-GD3 mouse antibody KM641 identified in the item 1 of Example 1, an FR amino acid sequence of VH of a human antibody was selected. Kabat *et al.* have classified the VH of various known human antibodies into three subgroups (HSG I to III) based on the homology of the amino acid sequences and then reported consensus sequences among respective subgroups (*Sequences of Proteins of Immunological Interest*). Since these consensus sequences have a possibility that the immunogenicity is reduced in human, it was decided to design a VH amino acid sequence of an anti-GD3 CDR-grafted antibody based on these consensus sequences. In order to produce an anti-GD3 CDR-grafted antibody having higher activity in designing it, it was decided to select an FR amino acid sequence having the highest homology with the FR amino acid sequence of VH of KM641, among FR amino acid sequences of consensus sequences of the three subgroups of VH of human antibodies. Results of the homology search are shown in Table 1. As shown in Table 1, the FR amino acid sequence of VH of KM641 showed the most high homology with the subgroup III.

(0111)

(Table 1)

Table 1
Homology between FR amino acid sequence of
consensus sequence of each subgroup of
human antibody H chain V region and
FR amino acid sequence of H chain V region of KM641

HSG I	HSG II	HSG III
62.1%	56.3%	78.2%

(0112)

Based on the above results, an amino acid sequence HV.0 of VH of the anti-GD3 CDR-grafted antibody was designed by grafting the CDR amino acid sequence of VH of the anti-GD3 mouse antibody KM641 to an appropriate position of the amino acid sequence of FR of the consensus sequence of subgroup III of VH of the human antibody.

Next, an amino acid sequence of VL of an anti-GD3 CDR-grafted antibody was designed as follows. In order to graft the CDR amino acid sequence of VL of the anti-GD3 mouse antibody KM641 identified in the item 1 of Example 1, an FR amino acid sequence of VL of a human antibody was selected. Kabat *et al.* have classified the VL of various known human antibodies into four subgroups (HSG I to IV) based on the homology of the amino acid sequences and then reported consensus sequences

among respective subgroups (*Sequences of Proteins of Immunological Interest*). Accordingly, similar to the H chain, an FR amino acid sequence having most high homology with the FR amino acid sequence of VL of KM641 was selected from FR amino acid sequences of consensus sequences of the four subgroups of VL of human antibodies. Results of the homology search are shown in Table 2. As shown in Table 2, the FR amino acid sequence of VL of KM641 showed the most high homology with the subgroup I.

(0113)

(Table 2)

Table 2
Homology between FR amino acid sequence of
consensus sequence of each subgroup of
human antibody L chain V region and
FR amino acid sequence of L chain V region of KM641

HSG I	HSG II	HSG III	HSG IV
76.2%	60.0%	62.5%	67.5%

(0114)

Based on the above results, an amino acid sequence LV.0 of VL of the anti-GD3 CDR-grafted antibody was designed by grafting the CDR amino acid sequence of VL of the anti-GD3 mouse antibody KM641 to an appropriate position of the amino acid sequence of FR of the consensus sequence of subgroup I of VL of the human antibody.

The thus designed amino acid sequence HV.0 of VH and amino acid sequence LV.0 of VL of the anti-GD3 CDR-grafted antibody are sequences in which only the CDR amino acid sequence of the anti-GD3 mouse antibody is grafted to the FR amino acid sequence of the selected human antibody. In general, in the human CDR-grafted antibodies, the antigen binding activity is reduced in many cases by grafting of a mouse antibody CDR amino acid sequence alone. Thus, in order to avoid the reduction of the antigen binding activity of antibodies, among FR amino acid residues different between a human antibody and a mouse antibody, amino acid residues considered to be exerting influences on the antigen binding activity are grafted together with a CDR amino acid sequence. Accordingly, identification of FR amino acid residues considered to have influences on the activity was carried out.

First, a three-dimensional structure of an antibody V region comprised of the amino acid sequence HV.0 of VH and amino acid sequence LV.0 of VL of anti-GD3 CDR-grafted antibody designed in the above (hereinafter referred to as "HV0LV0") was

constructed using computer-modeling. The production of three-dimensional structure coordinates was carried out using a software AbM (manufactured by Oxford Molecular) and the display of three-dimensional structure was carried out using a software Pro-Explore (manufactured by Oxford Molecular) in accordance with the respective manufacture's instructions. Also, a computer model of the three-dimensional structure of the V region of anti-GD3 mouse antibody KM641 was constructed in the same manner. In addition, a three-dimensional structure model of a modified HV0LV0 having an amino acid sequence in which at least one amino acid residue different from anti-GD3 mouse antibody KM641 in the FR amino acid sequences of VH and VL of HV0LV0 was substituted by the amino acid residues of positions corresponding to the anti-GD3 mouse antibody KM641 in order was also constructed. Three-dimensional structures of V regions of the anti-GD3 mouse antibody KM641, HV0LV0 and modified product were compared. As a result, amino acid residues considered to have influences on the antigen binding activity by changing three-dimensional structure of the antigen-binding region were selected from the FR amino acid residues of HV0LV0. As a result of substituting the thus selected FR amino acid residues of HV0LV0 by the amino acid residues found in the mouse antibody KM641, an amino acid sequence hKM641H of VH of the anti-GD3 CDR-grafted antibody represented by SEQ ID NO:9 and an amino acid sequence hKM641L of VL of the anti-GD3 CDR-grafted antibody represented by SEQ ID NO:10 were designed. In the hKM641H, the 10th position Gly, the 11th position Leu, the 20th position Leu, the 28th position Thr, the 84th position Asn, the 91st position Thr, the 95th position Tyr, the 97th position Ala and the 115th position Val in the FR amino acid sequence of HV.0 were replaced by Asp, Phe, Val, Ala, Arg, Ser, Phe, Thr and Leu, respectively, as amino acid residues of positions corresponding to the VH of anti-GD3 mouse antibody KM641. In the hKM641L, the 49th position Tyr, the 65th position Ser and the 71st position Phe in the FR amino acid sequence of LV.0 were replaced by Tyr, Ser and Phe, respectively, as amino acid residues of positions corresponding to the VL of anti-GD3 mouse antibody KM641.

(0115)

(2) Construction of cDNA encoding VH of anti-GD3 CDR-grafted antibody

A cDNA encoding the anti-GD3 CDR-grafted antibody VH amino acid sequence hKM641H designed in the item 3(1) of Example 1 was constructed as follows.

First, the designed amino acid sequence was ligated with the secretory signal sequence of H chain of anti-GD3 mouse antibody KM641 represented by SEQ ID NO:1 to produce a full antibody amino acid sequence. Next, the amino acid sequence is converted into genetic codons. When two or more genetic codons were present for

one amino acid residue, corresponding genetic codon was determined by taking the frequency of codon usage found in nucleotide sequences of antibody genes into consideration (*Sequences of Proteins of Immunological Interest*). A nucleotide sequence of cDNA encoding the complete antibody V region amino acid sequence was designed by ligating the thus determined genetic codons, and nucleotide sequences of primer binding site for use in the PCR amplification (including restriction enzyme recognition sequences for cloning into a humanized antibody expression vector) were added to the 5'-terminal and 3'-terminal. The thus designed nucleotide sequence was divided into a total of 6 nucleotide sequences from the 5'-terminal side, each having about 100 bases (adjoining nucleotide sequences are designed such that the termini have an overlapping sequence of about 20 bases), and they were synthesized in alternating order of the sense chain and the antisense chain using an automatic DNA synthesizer (380A, manufactured by Applied Biosystems) (0116)

Specifically, 6 synthetic DNA fragments of SEQ ID NO:11 to SEQ ID NO:16 were synthesized. Each DNA was added to 50 µl of a buffer comprising 10 mM Tris-HCl (pH 8.3), 50 mM potassium chloride, 1.5 mM magnesium chloride, 0.001% gelatin, 200 µM dNTPs, 0.5 µM M13 primer RV (manufactured by Takara Shuzo), 0.5 µM M13 primer M4 (manufactured by Takara Shuzo) and 2 units of TaKaRa Taq DNA polymerase (manufactured by Takara Shuzo) to give a final concentration of 0.1 µM, and the solution was covered with 50 µl of mineral oil and set to a DNA thermal cycler (PJ480, manufactured by PERKIN ELMER) to carry out 30 cycles of the reaction, each cycle including 2 minutes at 94°C, 2 minutes at 55°C and 2 minutes at 72°C. The reaction solution was purified using QIAquick PCR Purification Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions and made into 30 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT, 100 µg/ml BSA and 0.01% Triton X-100, and 10 units of a restriction enzyme *NorI* (manufactured by Takara Shuzo) were further added to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 0.2 µg of an *ApaI-NorI* fragment of about 0.44 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0117)

Next, 3 µg of a plasmid pBluescript SK(-) was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol and added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT, 100 µg/ml BSA and 0.01% Triton X-100, and 10 units of a restriction enzyme *NotI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 2 µg of an *ApaI-NotI* fragment of about 2.95 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0118)

Next, 0.1 µg of the *ApaI-NotI* fragment of the PCR product of VH of anti-GD3 CDR-grafted antibody and 0.1 µg of the *ApaI-NotI* fragment of the plasmid pBluescript SK(-), both obtained in the above, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained recombinant plasmid DNA solution, an *Escherichia coli* line HB101 was transformed. Each plasmid DNA was produced from 10 clones of the transformants, allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence, and as a result, the plasmid pHKM641H shown in Fig. 1 having the nucleotide sequence of interest was obtained.

(0119)

(3) Construction of cDNA encoding VL of anti-GD3 CDR-grafted antibody

A cDNA encoding the anti-GD3 CDR-grafted antibody VL amino acid sequence hKM641L designed in the item 3(1) of Example 1 was constructed as follows using the PCR similar to the VH. In this case, the amino acid sequence of L chain of anti-GD3 mouse antibody KM641 represented by SEQ ID NO:2 was used as the secretory signal sequence.

(0120)

First, 6 synthetic DNA fragments of SEQ ID NO:17 to SEQ ID NO:22 were synthesized using an automatic DNA synthesizer (380A, manufactured by Applied Biosystems). Each of the thus synthesized DNA fragments was added to 50 µl of a

buffer comprising 10 mM Tris-HCl (pH 8.3), 50 mM potassium chloride, 1.5 mM magnesium chloride, 0.001% gelatin, 200 μ M dNTPs, 0.5 μ M M13 primer RV (manufactured by Takara Shuzo), 0.5 μ M M13 primer M4 (manufactured by Takara Shuzo) and 2 units of TaKaRa Taq DNA polymerase (manufactured by Takara Shuzo) to give a final concentration of 0.1 μ M, and the solution was covered with 50 μ l of mineral oil and set to a DNA thermal cycler (PJ480, manufactured by PERKIN ELMER) to carry out 30 cycles of the reaction, each cycle including 2 minutes at 94°C, 2 minutes at 55°C and 2 minutes at 72°C. The reaction solution was purified using QIAquick PCR Purification Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions and made into 30 μ l of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 μ g/ml BSA, and 10 units of a restriction enzyme *Eco*RI (manufactured by Takara Shuzo) and a restriction enzyme *Sp*II (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 0.2 μ g of an *Eco*RI-*Sp*II fragment of about 0.39 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0121)

Next, 3 μ g of a plasmid pBSL3 described in Japanese Published Unexamined Patent Application No. 257893/98 was added to 10 μ l of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 μ g/ml BSA, and 10 units of a restriction enzyme *Eco*RI (manufactured by Takara Shuzo) and a restriction enzyme *Sp*II (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 2 μ g of an *Eco*RI-*Sp*II fragment of about 2.95 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0122)

Next, 0.1 μ g of the *Eco*RI-*Sp*II fragment of the PCR product of VL of anti-GD3 CDR-grafted antibody and 0.1 μ g of the *Eco*RI-*Sp*II fragment of the plasmid pBSL3, both obtained in the above, were added to 20 μ l in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained recombinant plasmid DNA solution, an *E. coli* line HB101 was transformed. Each plasmid DNA was prepared from 10 clones of the transformants, allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to electrophoresis

using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence, and as a result, the plasmid pKM641L shown in Fig. 2 having the nucleotide sequence of interest was obtained.

(0123)

5 4. Activity evaluation of anti-GD3 CDR-grafted antibody by transient expression using animal cell

In order to carry out activity evaluation of anti-GD3 CDR-grafted antibody more quickly, transient expression of anti-GD3 CDR-grafted antibody was carried out as follows using COS-7 cell (ATCC CRL 1651).

10 (1) Construction of anti-GD3 chimeric antibody KM871 transient expression vector pT641

In order to use as a positive control in the activity evaluation of anti-GD3 CDR-grafted antibody by transient expression, a transient expression vector pT641 for anti-GD3 chimeric antibody KM871 was constructed as follows.

15 (0124)

Since the efficiency of transient expression using animal cells generally depends on the copy number of introduced expression vector, it was considered that smaller expression vector has higher expression efficiency. Accordingly, the transient expression vector pT641 for anti-GD3 chimeric antibody KM871 was constructed as follows using the transient expression vector pT796 for anti-ganglioside GM2 chimeric antibody KM966 described in Japanese Published Unexamined Patent Application No. 257893/98 and the plasmid pKM641HF1 having VH of anti-GD3 mouse antibody KM641 and plasmid pKM641LA2 having VL of anti-GD3 mouse antibody KM641 described in Japanese Published Unexamined Patent Application No. 304989/93.

25 (0125)

First, 3 µg of the plasmid pKM641LA2 having VL of anti-GD3 mouse antibody KM641 described in Japanese Published Unexamined Patent Application No. 304989/93 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 50 mM sodium chloride, 10 mM magnesium chloride and 1 mM DTT and the solution was further mixed with 10 units of a restriction enzyme *Hind*III (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol and added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride and 1 mM DTT, and 10 units of a restriction enzyme *Eco*RI (manufactured by Takara Shuzo) were further added to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 0.2 µg of an *Hind*III-*Eco*RI

fragment of about 0.35 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0126)

Next, 3 µg of the transient expression vector pT796 for anti-ganglioside GM2 chimeric antibody KM966 described in Japanese Published Unexamined Patent Application No. 257893/98 was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 µg/ml BSA, and the solution was further mixed with 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) and a restriction enzyme *SpII* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 2 µg of an *EcoRI-SpII* fragment of about 9.20 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0127)

Next, synthetic DNA fragments respectively having the nucleotide sequences represented by SEQ ID NO:23 and SEQ ID NO:24 were synthesized using an automatic DNA synthesizer (380A, manufactured by Applied Biosystems). A 0.3 µg of each of the thus obtained synthetic DNA fragments was added to 15 µl of sterile water and heated at 65°C for 5 minutes. After allowing the reaction solution to stand at room temperature for 30 minutes, this was mixed with 2 µl of a 10 × buffer solution (500 mM Tris-HCl (pH 7.6), 100 mM magnesium chloride, 50 mM DTT) and 2 µl of 10 mM ATP and further mixed with 10 units of T4 Polynucleotide Kinase (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 30 minutes to thereby phosphorylate the 5'-terminal.

(0128)

Next, 0.1 µg of the *HindIII-EcoRI* fragment derived from the plasmid pKM641LA2, 0.1 µg of the *EcoRI-SpII* fragment derived from the plasmid pT796 and 0.05 µg of the phosphorylated synthetic DNA, obtained in the above, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained recombinant plasmid DNA solution, an *E. coli* HB101 was transformed to obtain the plasmid pT796H641L shown in Fig. 3. The thus obtained plasmid (10 µg) was allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to an electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence, and as a result, it was confirmed that a plasmid into which the DNA of interest was cloned was obtained.

(0129)

Next, 3 µg of a plasmid pBluescript SK(-) (manufactured by Stratagene) was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 50 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 µg/ml BSA and the solution was further mixed with 10 units of a restriction enzyme *Xba*I (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 30 µl of a buffer comprising 30 mM sodium acetate (pH 5.0), 100 mM sodium chloride, 1 mM zinc acetate and 5% glycerol, and further mixed with 30 units of a modification enzyme Mung Bean Nuclease (manufactured by Takara Shuzo) to carry out the reaction at 25°C for 15 minutes. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and further mixed with 10 units of a restriction enzyme *Apa*I (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of a Blunt end-*Apa*I fragment of about 2.95 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0130)

Next, 3 µg of the plasmid pKM641HF1 was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed with 10 units of a restriction enzyme *Eco*RI (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, and the 5' protruding end formed by the restriction enzyme digestion was changed to blunt end using DNA Blunting Kit (manufactured by Takara Shuzo). The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and further mixed with 10 units of a restriction enzyme *Apa*I (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 0.2 µg of a Blunt end-*Apa*I fragment of about 0.44 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0131)

Next, 0.1 µg of Blunt end-*Apa*I fragment derived from the plasmid pBluescript SK(-) and 0.1 µg of the Blunt end-*Apa*I fragment derived from the plasmid pKM641HF1, obtained in the above, were added to 20 µl in total volume of sterile

water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained recombinant plasmid DNA solution, an *E. coli* HB101 was transformed to obtain the plasmid pBS641H shown in Fig. 4.

(0132)

5 Next, 3 µg of the plasmid pT796H641L obtained in the above was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT and the solution was further mixed with 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer
10 comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT, 100 µg/ml BSA and 0.01% Triton X-100, and further mixed with 10 units of a restriction enzyme *NotI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 0.2 µg of an *ApaI-NotI* fragment of about 9.16 kb using
15 QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0133)

 Next, 3 µg of the plasmid pBS641H obtained in the above was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1
20 mM DTT and the solution was further mixed with 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT, 100 µg/ml BSA and 0.01% Triton X-100, and further mixed with 10 units of a
25 restriction enzyme *NotI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an *ApaI-NotI* fragment of about 0.44 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

30 (0134)

 Next, 0.1 µg of the *ApaI-NotI* fragment derived from the plasmid pT796H641L and 0.1 µg of the *ApaI-NotI* fragment derived from the plasmid pBS641H, obtained in the above, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus
35 obtained recombinant plasmid DNA solution, an *E. coli* HB101 was transformed to obtain the plasmid pT641 shown in Fig. 5.

(0135)

(2) Construction of an anti-GD3 CDR-grafted antibody KM871 transient expression vector

A transient expression vector for anti-GD3 CDR-grafted antibody was constructed as follows using the transient expression vector pT641 for anti-GD3 chimeric antibody described in the item 4(1) of Example 1 and the plasmids phKM641H and phKM641L described in the items 3(2) and(3) of Example 1.

(0136)

First, 3 µg of the plasmid phKM641H obtained in the item 3(2) of Example 1 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT and the solution was further mixed with 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT, 100 µg/ml BSA and 0.01% Triton X-100, and further mixed with 10 units of a restriction enzyme *NotI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 0.2 µg of an *ApaI-NotI* fragment of about 0.44 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0137)

Next, 3 µg of the plasmid pT641 obtained in the item 4(1) of Example 1 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed with 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT, 100 µg/ml BSA and 0.01% Triton X-100, and further mixed with 10 units of a restriction enzyme *NotI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an *ApaI-NotI* fragment of about 9.16 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0138)

Next, 0.1 µg of the *ApaI-NotI* fragment derived from the plasmid phKM641H and 0.1 µg of the *ApaI-NotI* fragment derived from the plasmid pT641

were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained recombinant plasmid DNA solution, an *E. coli* HB101 was transformed to obtain the plasmid pT641HCDR shown in Fig. 6.

5 (0139)

Next, 3 µg of the plasmid pHKM641L obtained in the item 3(3) of Example 1 was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 µg/ml BSA, and the solution was further mixed with 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) and a restriction enzyme *SpI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 0.2 µg of an *EcoRI-SpI* fragment of about 0.39 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

15 (0140)

Next, 3 µg of each of the plasmid pT641 obtained in the item 4(1) of Example 1 and the plasmid pT641HCDR obtained in the above was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 µg/ml BSA, and the solution was further mixed with 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) and a restriction enzyme *SpI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an *EcoRI-SpI* fragment of about 9.20 kb derived from each plasmid using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

25 (0141)

Next, 0.1 µg of the *EcoRI-SpI* fragment derived from the plasmid pHKM641L and 0.1 µg of the *EcoRI-SpI* fragment derived from the plasmid pT641, obtained in the above, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Also, 0.1 µg of the *EcoRI-SpI* fragment derived from the plasmid pHKM641L and 0.1 µg of the *EcoRI-SpI* fragment derived from the plasmid pT641HCDR, obtained in the above, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using each of the thus obtained recombinant plasmid DNA solutions, an *E. coli* HB101 was transformed to obtain the plasmids pT641LCDR and pT641HLCDR shown in Fig. 7.

(0142)

(3) Activity evaluation of anti-GD3 chimeric antibody and anti-GD3 CDR-grafted antibody by transient expression using animal cells

Transient expression of antibodies was carried out using the transient expression vector pT641 for anti-GD3 chimeric antibody, the transient expression vector pT641HLCDR for anti-GD3 CDR-grafted antibody and the transient expression vectors pT641HCDR and pT641LCDR for anti-GD3 hybrid antibodies having V regions of mouse antibody and human CDR-grafted antibody, obtained in the items 4(1) and (2) of Example 1.

(0143)

COS-7 cells (ATCC CRL 1651) were dispensed in 2 ml at a density of 1×10^5 cells/ml into a 6 well plate (manufactured by Falcon) and cultured overnight at 37°C. By adding 2 µg of each expression vector to 100 µl of OPTI-MEM medium (manufactured by GIBCO BRL) and further adding a solution in which 10 µl of LIPOFECTAMINE Reagent (manufactured by GIBCO BRL) was added to 100 µl of the OPTI-MEM medium, the reaction was carried out at room temperature for 40 minutes to effect formation of a DNA-liposome complex. The overnight-cultured COS-7 cells were washed twice with 2 ml of the OPTI-MEM medium (manufactured by GIBCO BRL), mixed with a solution prepared by adding 0.8 ml of OPTI-MEM medium to the complex-containing solution and cultured at 37°C for 7 hours, the solution was discarded and then the cells were mixed with 2 ml of 10% FBS-containing DME medium (manufactured by GIBCO BRL) and cultured at 37°C. After the introduction of each expression vector, the culture supernatant was recovered 72 hours thereafter and its concentration operation was carried out as occasion demands, and then binding activity of the anti-GD3 humanized antibody for GD3 (manufactured by DIA-IATRON) in the culture supernatant was measured by the ELISA method described in the item 2 of Example 1, by measuring concentration of the anti-GD3 humanized antibody in the culture supernatant by the ELISA method described in the item 4 which will be described later and calculating the activity from the measured values as relative activity (%) when the activity of the positive control anti-GD3 chimeric antibody is defined as 100. Results are shown in Fig. 8. As shown in Fig. 8, the anti-GD3 hybrid antibody derived from the transient expression vector pT641HCDR (VH is derived from a human CDR-grafted antibody and VL is derived from a chimeric antibody) showed almost the same binding activity of that of the anti-GD3 hybrid antibody derived from the transient expression vector pT641, but the anti-GD3 hybrid antibody derived from the transient expression vector pT641LCDR (VH is derived from a chimeric antibody and VL is

derived from a human CDR-grafted antibody) showed a binding activity of about 50% of the anti-GD3 chimeric antibody. In addition, the anti-GD3 CDR-grafted antibody derived from the transient expression vector pT641HLCADR showed a binding activity of only about 5% of the anti-GD3 chimeric antibody, showing considerable decrease in the activity.

(0144)

Based on these results, binding activity of the anti-GD3 CDR-grafted antibody having VH and VL designed in the item 3(1) of Example 1 is considerably reduced and the main cause of the activity reduction is considered to be due to VL, but since about 50% of the activity is found by its combination with the VH of anti-GD3 chimeric antibody, it was suggested that there is a problem particularly at the interaction region with VH. Accordingly, an attempt was made to increase the antibody activity by further modifying amino acid residues of the VL designed in the item 3(1) of Example 1.

(0145)

(4) Measurement of the concentration of humanized antibody in transient expression culture supernatant by ELISA method

A solution (50 μ l) prepared by diluting a goat anti-human IgG (γ -chain) antibody (manufactured by Medical & Biological Laboratories) 400 folds with PBS was dispensed in each well of a 96 well plate for ELISA use (manufactured by Greiner) and allowed to react at 4°C overnight. After discarding the antibody solution, 1% BSA-PBS was added in 100 μ l/well and allowed to react at room temperature for 1 hour to thereby block the remaining active groups. After discarding 1% BSA-PBS, the transient expression culture supernatants obtained in the item 4(3) of Example 1 or diluted solutions of the purified anti-GD3 chimeric antibody KM871 were added in 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction, each well was washed with Tween-PBS and then a solution prepared by diluting a peroxidase-labeled mouse anti-human κ L chain antibody (manufactured by Zymed) 500 folds with PBS was added in 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, an ABTS substrate solution (a solution prepared by dissolving 0.55 g of 2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid) in 1 L of 0.1 M citrate buffer (pH 4.2) and adding 1 μ l/ml of hydrogen peroxide just before the use) was added in 50 μ l/well for color development and OD415 was measured.

(0146)

5. Increase of the binding activity by modifying amino acid residues of VL of anti-GD3 CDR-grafted antibody

Increase of the binding activity by modifying amino acid residues of VL of the anti-GD3 CDR-grafted antibody designed in the item 3(1) of Example 1 was carried out as follows.

(1) Modification of amino acid residues of VL of anti-GD3 CDR-grafted antibody and construction of cDNA encoding the modified VL

First, amino acid residues which are amino acid residues positioned in the interaction region of VH and VL and amino acid residues considered to be exerting influence on the three-dimensional structure of each CDR of VL and are also different from the amino acid residues of mouse antibody in the VL of human CDR-grafted antibody were identified from computer models of the three-dimensional structures of various antibodies constructed in the item 3(1) of Example 1. As a result, the 7th position Ser, the 8th position Pro, the 12th position Ser, the 41st position Gly, the 44th position Pro, the 72nd position Thr, the 77th position Ser, the 83rd position Phe and the 87th Tyr of hKM641L as the amino acid sequence of VL of anti-GD3 CDR-grafted antibody represented by SEQ ID NO:10 were identified. By replacing these amino acid residues by amino acid residues found in the mouse antibody, 8 VLs of modified anti-GD3 CDR-grafted antibody were designed. That is, the 8th position Pro, the 12th position Ser, the 44th position Pro and the 87th position Tyr among the hKM641L amino sequence were respectively replaced by Ala, Pro, Val and Phe in the case of hKM641NL, and the 7th position Ser, the 8th position Pro and the 12th position Ser among the hKM641L amino sequence respectively by Thr, Ala and Pro in the case of hKM641Lm-1, the 87th position Tyr among the hKM641L amino sequence by Phe in the case of hKM641Lm-4, the 41st position Gly and the 44th position Pro among the hKM641L amino sequence respectively by Asp and Val in the case of hKM641Lm-6, the 72nd position Thr, the 77th position Ser and the 83rd position Phe among the hKM641L amino sequence respectively by Ser, Asn and Ile in the case of hKM641Lm-7, the 77th position Ser among the hKM641L amino sequence by Asn in the case of hKM641Lm-8, the 83rd position Phe and the 87th position Tyr among the hKM641L amino sequence respectively by Ile and Phe in the case of hKM641Lm-9, and the 41st position Gly, the 44th position Pro and the 83rd position Phe among the hKM641L amino sequence respectively by Asp, Val and Ile in the case of hKM641Lm-69. Among these modified VLs, cDNA encoding each of 6 modified VLs excluding hKM641NL and hKM641Lm-69 was constructed as follows by PCR-aided mutagenesis. That is, antisense chain and sense chain DNA primers were synthesized using an

automatic DNA synthesizer (380A, manufactured by Applied Biosystems) for introducing mutation, and a first PCR was carried out in accordance with the method described in the item 3(2) of Example 1 using 1 ng of the plasmid phKM641L as the template and 0.5 μ M in final concentration of M13 primer RV (manufactured by Takara Shuzo) and the antisense chain DNA primer and M13 primer M4 (manufactured by Takara Shuzo) and the sense chain DNA primer. Each of the reaction solutions was purified using QIAquick PCR Purification Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions by eluting with 20 μ l of 10 mM Tris-HCl (pH 8.0), and then a second PCR was carried out using 5 μ l of each eluate in accordance with the method described in the item 3(2) of Example 1. The reaction solution was purified using QIAquick PCR Purification Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions and made into 30 μ l of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 μ g/ml BSA, and 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) and a restriction enzyme *SpI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 0.2 μ g of an *EcoRI-SpI* fragment of about 0.39 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0147)

Next, 0.1 μ g of the *EcoRI-SpI* fragment of the PCR product of VL of the modified anti-GD3 CDR-grafted antibody obtained in the above and 0.1 μ g of the *EcoRI-SpI* fragment of the plasmid pBSL3 obtained in the item 3(3) of Example 1 were added to 20 μ l in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained recombinant plasmid DNA solution, an *E. coli* HB101 was transformed. Each plasmid DNA was produced from 10 clones of the transformants, allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence to thereby obtain a plasmid having cDNA to which the intended modification was applied.

(0148)

Specifically, a plasmid phKM641Lm-1 having the nucleotide sequence represented by SEQ ID NO:27 was obtained by carrying out a series of the above operations using the synthetic DNA of SEQ ID NO:25 as the antisense chain DNA

primer and the synthetic DNA of SEQ ID NO:26 as the sense chain DNA primer. The hKM641Lm-1 as an amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:27.

(0149)

5 A plasmid phKM641Lm-4 having the nucleotide sequence represented by SEQ ID NO:30 was obtained by carrying out a series of the above operations using the synthetic DNA of SEQ ID NO:28 as the antisense chain DNA primer and the synthetic DNA of SEQ ID NO:29 as the sense chain DNA primer. The hKM641Lm-4 as an amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID
10 NO:30.

A plasmid phKM641Lm-6 having the nucleotide sequence represented by SEQ ID NO:33 was obtained by carrying out a series of the above operations using the synthetic DNA of SEQ ID NO:31 as the antisense chain DNA primer and the synthetic DNA of SEQ ID NO:32 as the sense chain DNA primer. The hKM641Lm-6 as an
15 amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:33.

(0150)

A plasmid phKM641Lm-7 having the nucleotide sequence represented by SEQ ID NO:36 was obtained by carrying out a series of the above operations using the
20 synthetic DNA of SEQ ID NO:34 as the antisense chain DNA primer and the synthetic DNA of SEQ ID NO:35 as the sense chain DNA primer. The hKM641Lm-7 as an amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:36.

A plasmid phKM641Lm-8 having the nucleotide sequence represented by
25 SEQ ID NO:39 was obtained by carrying out a series of the above operations using the synthetic DNA of SEQ ID NO:37 as the antisense chain DNA primer and the synthetic DNA of SEQ ID NO:38 as the sense chain DNA primer. The hKM641Lm-8 as an amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:39.

30 (0151)

A plasmid phKM641Lm-9 having the nucleotide sequence represented by
35 SEQ ID NO:42 was obtained by carrying out a series of the above operations using the synthetic DNA of SEQ ID NO:40 as the antisense chain DNA primer and the synthetic DNA of SEQ ID NO:41 as the sense chain DNA primer. The hKM641Lm-9 as an amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:42.

Among the modified VLs, cDNA encoding the hKM641NL was constructed by synthesizing 6 synthetic DNA fragments of SEQ ID NOs:17, 22 and 43 to 46 using an automatic DNA synthesizer (380A, manufactured by Applied Biosystems) and carrying out the procedure described in the item 3(3) of Example 1 using these fragments. As a result, a plasmid phKM641NL having the nucleotide sequence represented by SEQ ID NO:47 in which the modification of interest was effected was obtained. The hKM641NL as an amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:47.

(0152)

Among the modified VLs, cDNA encoding the hKM641Lm-69 was constructed as follows using the plasmids phKM641Lm-6 and phKM641Lm-9 having cDNA of modified VL obtained in the above.

First, 3 µg of the plasmid phKM641Lm-6 was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM of sodium chloride, 10 mM magnesium chloride and 1 mM DTT and 10 units of restriction enzymes *EcoRI* (manufactured by Takara Shuzo) and *PstI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 0.3 µg of an *EcoRI-PstI* fragment of about 0.30 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0153)

Next, 3 µg of the plasmid phKM641Lm-9 was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM of sodium chloride, 10 mM magnesium chloride and 1 mM DTT and 10 units of restriction enzymes *EcoRI* (manufactured by Takara Shuzo) and *PstI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 2 µg of an *EcoRI-PstI* fragment of about 3.05 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0154)

Next, 0.1 µg of the *EcoRI-PstI* fragment derived from the plasmid phKM641Lm-6 and 0.1 µg of the *EcoRI-PstI* fragment derived from the plasmid phKM641Lm-9, both obtained in the above, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained recombinant plasmid DNA solution, an *E. coli* HB101 was transformed to obtain the plasmid phKM641Lm-69 shown in Fig. 9. The

reaction was carried out using 10 µg of the thus obtained plasmid in accordance with the manufacturer's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then the resulting mixture was subjected to electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence, and as a result, it was confirmed that it has the nucleotide sequence represented by SEQ ID NO:48 in which the modification of interest was carried out. The hKM641Lm-69 as an amino acid sequence encoded by the nucleotide sequence is also represented by SEQ ID NO:48.

(0155)

(2) Construction of transient expression vector for anti-GD3 CDR-grafted antibodies having modified VL

Transient expression vectors for anti-GD3 CDR-grafted antibodies having various modified VLs were constructed as follows using the plasmids having cDNA molecules encoding various modified VLs, obtained in the item 5(1) of Example 1, and the transient expression vector pT641HCDR for anti-GD3 hybrid antibodies obtained in the item 4(2) of Example 1.

(0156)

First, 3 µg of each of the plasmids phKM641NL, phKM641Lm-1, phKM641Lm-4, phKM641Lm-6, phKM641Lm-7, phKM641Lm-8, phKM641Lm-9 and phKM641Lm-69 obtained in the item 5(1) of Example 1 was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM of sodium chloride, 10 mM magnesium chloride, 1 mM DTT and 100 µg/ml of BSA, and 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) and a restriction enzyme *SpI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 0.2 µg of an *EcoRI-SpI* fragment of about 0.39 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacturer's instructions.

(0157)

Next, 0.1 µg of each of the thus obtained *EcoRI-SpI* fragments of various modified VLs and 0.1 µg of the *EcoRI-SpI* fragment of the transient expression vector pT641HCDR for anti-GD3 hybrid antibody obtained in the item 4(2) of Example 1, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using the thus obtained respective recombinant plasmid DNA solutions, an *E. coli* HB101 was transformed to obtain the transient expression vectors for anti-GD3 CDR-grafted antibodies having various modified VLs, pT641HLCDRNL, pT641HLCDRLm-1, pT641HLCDRLm-4,

pT641HLCDRm-6, pT641HLCDRm-7, pT641HLCDRm-8, pT641HLCDRm-9 and pT641HLCDRm-69, shown in Fig. 10.

(0158)

(3) Evaluation of activity of anti-GD3 CDR-grafted antibody having modified VLs by transient expression using animal cell

Transient expression and evaluation of activity of each antibody were carried out in accordance with the method described in the item 4(3) of Example 1 using the transient expression vector pT641 for anti-GD3 chimeric antibody and the transient expression vector pT641HLCDR for anti-GD3 CDR-grafted antibody, both obtained in the item 4(2) of Example 1, and the transient expression vectors, pT641HLCDRNL, pT641HLCDRm-1, pT641HLCDRm-4, pT641HLCDRm-6, pT641HLCDRm-7, pT641HLCDRm-8, pT641HLCDRm-9 and pT641HLCDRm-69, for anti-GD3 CDR-grafted antibodies having various modified VLs, obtained in the item 5(2) of Example 1. Results are shown in Fig. 11. As shown in Fig. 11, the modified anti-GD3 CDR-grafted antibodies derived from the transient expression vectors, pT641HLCDRm-6, pT641HLCDRm-7, pT641HLCDRm-9 and pT641HLCDRm-69, showed increased binding activity in comparison with the anti-GD3 CDR-grafted antibodies before modification, particularly, the modified anti-GD3 CDR-grafted antibody derived from the transient expression vector pT641HLCDRm-69 showed its binding activity of about 80% of the anti-GD3 chimeric antibody. On the other hand, the modified anti-GD3 CDR-grafted antibodies derived from other transient expression vectors showed no increase in the binding activity in comparison with the anti-GD3 CDR-grafted antibodies before modification. Based on these results, it was found that, among the modified amino acid residues of VL identified in the item 5(1) of Example 1, the 41st position, 44th position, 83rd position and 87th position amino acid residues greatly contribute to the increase of binding activity. It was also found that simultaneous substitution of two amino acid residues of the 41st position and 44th position or the 83rd position and 87th position contributes to synergistic increase in the activity, and simultaneous modification of four amino acid residues of the 41st position, 44th position, 83rd position and 87th position also contributes to synergistic increase in the activity. Based on the computer models of three-dimensional structures of V regions of various antibodies constructed in the item 3(1) of Example 1, it was suggested that the 41st position and 44th position amino acid residues exert influence on the interaction with VH, and the 83rd position and 87th position amino acid residues on the three-dimensional structure of CDR3 of VL, and it was considered that three-dimensional structure of an antibody as a whole is suitably maintained by modifying

these amino acid residues into amino acid residues found in a mouse antibody and, as a result, its binding activity increases. This information shows that it is necessary to take the interaction regions of VH and VL also into consideration in preparing human CDR-grafted antibodies and that it is necessary to carry out various examinations on the identification of these interaction regions based on the information from three-dimensional structures of antibody V regions. With regard to these interaction regions, it can be easily analogize that they are different in individual antibodies and it is difficult to find a general law at present, so that it is necessary to carry out trial and error in response to each antibody of interest.

(0159)

6. Stable expression of anti-GD3 CDR-grafted antibody using animal cell

Based on the results described in the item 5(3) of Example 1, it was suggested that the anti-GD3 CDR-grafted antibodies derived from transient expression vectors pT641HLCDR_Lm-9 and pT641HLCDR_Lm-69 have about 20% and about 80% of binding activity for GD3, respectively, in comparison with anti-GD3 chimeric antibodies. Accordingly, in order to evaluate the activity of these anti-GD3 CDR-grafted antibodies more in detail, transformed cell lines capable of stably expressing anti-GD3 CDR-grafted antibodies were obtained by the following method, and purification of anti-GD3 CDR-grafted antibodies from culture supernatants of the transformed cell lines was carried out. Also, for comparison, stable expression and purification of the anti-GD3 CDR-grafted antibody derived from the transient expression vector pT641HLCDR were carried out in the same manner.

(0160)

(1) Construction of stable expression vector for anti-GD3 CDR-grafted antibody

Stable expression vectors were constructed by introducing a resistance gene against an agent G418 and the dhfr gene into various transient expression vectors constructed in the item 5(2) of Example 1.

(0161)

First, 3 μ g of the humanized antibody expression vector pKANTEX93 described in WO 97/10354 was added to 10 μ l of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and 10 units of a restriction enzyme *Bam*HI (manufactured by Takara Shuzo) and a restriction enzyme *Xho*I (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 1 μ g of a *Bam*HI-*Xho*I fragment of about

these amino acid residues into amino acid residues found in a mouse antibody and, as a result, its binding activity increases. This information shows that it is necessary to take the interaction regions of VH and VL also into consideration in preparing human CDR-grafted antibodies and that it is necessary to carry out various examinations on the identification of these interaction regions based on the information from three-dimensional structures of antibody V regions. With regard to these interaction regions, it can be easily analogized that they are different in individual antibodies and it is difficult to find a general law at present, so that it is necessary to carry out trial and error in response to each antibody of interest.

(0159)

6. Stable expression of anti-GD3 CDR-grafted antibody using animal cell

Based on the results described in the item 5(3) of Example 1, it was suggested that the anti-GD3 CDR-grafted antibodies derived from transient expression vectors pT641HLCDR_Lm-9 and pT641HLCDR_Lm-69 have about 20% and about 80% of binding activity for GD3, respectively, in comparison with anti-GD3 chimeric antibodies. Accordingly, in order to evaluate the activity of these anti-GD3 CDR-grafted antibodies more in detail, transformed cell lines capable of stably expressing anti-GD3 CDR-grafted antibodies were obtained by the following method, and purification of anti-GD3 CDR-grafted antibodies from culture supernatants of the transformed cell lines was carried out. Also, for comparison, stable expression and purification of the anti-GD3 CDR-grafted antibody derived from the transient expression vector pT641HLCDR were carried out in the same manner.

(0160)

(1) Construction of stable expression vector for anti-GD3 CDR-grafted antibody

Stable expression vectors were constructed by introducing a resistance gene against an agent G418 and the dhfr gene into various transient expression vectors constructed in the item 5(2) of Example 1.

(0161)

First, 3 μ g of the humanized antibody expression vector pKANTEX93 described in WO 97/10354 was added to 10 μ l of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and 10 units of a restriction enzyme *Bam*HI (manufactured by Takara Shuzo) and a restriction enzyme *Xho*I (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 1 μ g of a *Bam*HI-*Xho*I fragment of about

8.68 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0162)

Next, 3 µg of each of the transient expression vectors, pT641HLCDRm-9, pT641HLCDRm-69 and pT641HLCDR, was added to 10 µl of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and 10 units of a restriction enzyme *Bam*HI (manufactured by Takara Shuzo), a restriction enzyme *Xho*I (manufactured by Takara Shuzo) and a restriction enzyme *Sst*I (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 1 µg of a *Bam*HI-*Xho*I fragment of about 4.90 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0163)

Next, 0.1 µg of the *Bam*HI-*Xho*I fragment derived from the plasmid pKANTEX93 and 0.1 µg of the *Bam*HI-*Xho*I fragment derived from each of the transient expression vectors, obtained in the above, were added to 20 µl in total volume of sterile water and ligated using Ready-To-Go T4 DNA Ligase (manufactured by Pharmacia). Using each of the thus obtained recombinant plasmid DNA solutions, an *E. coli* HB101 was transformed to obtain the anti-GD3 CDR-grafted antibody stable expression vectors, pKANTEX641HLCDRm-9, pKANTEX641HLCDRm-69 and pKANTEX641HLCDR, shown in Fig. 12.

(0164)

(2) Stable expression of anti-GD3 CDR-grafted antibody using animal cell

Using the various stable expression vectors for anti-GD3 CDR-grafted antibody obtained in the item 6(1) of Example 1, expression of anti-GD3 CDR-grafted antibody in animal cells was carried out as follows.

Each of the anti-GD3 CDR-grafted antibody expression vectors (4 µg) was introduced into 4×10^6 cells of a rat myeloma cell line YB2/0 cell (ATCC CRL 1662) by the electroporation method (*Cytotechnology*, 3, 133 (1990)), and the cells were suspended in 40 ml of RPMI640-FBS(19) (RPMI640 medium containing 10% of fetal bovine serum (FBS)) and dispensed in 200 µl/well into a 96 well microtiter plate (manufactured by Sumitomo Bakelite). After culturing at 37°C for 24 hours in a 5% CO₂ incubator, G418 was added to give a concentration of 0.5 mg/ml, followed by culturing for 1 to 2 weeks. Culture supernatants were recovered from wells in which colonies of transformants showing G418-resistance were formed and became confluent,

and the antigen binding activity of various anti-GD3 CDR-grafted antibodies in the supernatants was measured by the ELISA shown in the item 2 of Example 1 (a peroxidase-labeled goat anti-human IgG(γ) antibody was used as the secondary antibody).

5 (0165)

With regard to the transformants of wells in which expression of anti-GD3 CDR-grafted antibodies was found in the culture supernatants, in order to increase antibody expression quantity making use of a dhfr gene amplification system, they were suspended in the RPMI1640-FBS(10) medium containing 0.5 mg/ml of G418 and 50 nM methotrexate (hereinafter referred to as "MTX": manufactured by SIGMA) as an inhibitor of the dhfr gene product dihydrofolate reductase (hereinafter referred to as "DHFR") to give a density of 1 to 2×10^5 cells/ml, and dispensed in 2 ml into a 24 well plate (manufactured by Greiner). By culturing at 37°C for 1 to 2 weeks in a 5% CO₂ incubator, transformants showing 50 nM MTX resistance were induced. When the transformants became confluent, the antigen binding activity of various anti-GD3 CDR-grafted antibodies in the supernatants was measured by the ELISA shown in the item 2 of Example 1. With regard to the transformants of wells in which expression of anti-GD3 CDR-grafted antibodies was found in the culture supernatants, the MTX concentration was increased to 100 nM and then to 200 nM by the method similar to the above, and transformants capable of growing in the RPMI1640-FBS(10) medium containing 0.5 mg/ml of G418 and 200 nM of MTX and of highly expressing anti-GD3 CDR-grafted antibodies were finally obtained. With regard to the thus obtained transformants, single cell isolation (cloning) was carried out twice by limiting dilution analysis. The transformed cell clones obtained by introducing the stable expression vectors, pKANTEX641HLCDR_Lm-9, pKANTEX641HLCDR_Lm-69 and pKANTEX641HLCDR, were named KM8870, KM8871 and KM8869, respectively, and expression levels of anti-GD3 CDR-grafted antibody by transformed cell clones were about 5 $\mu\text{g}/10^6$ cells/24 hr, about 10 $\mu\text{g}/10^6$ cells/24 hr and about 30 $\mu\text{g}/10^6$ cells/24 hr, respectively.

30 (0166)

(3) Purification of anti-GD3 CDR-grafted antibody from culture filtrate

Each of the transformed cell clones, KM8870, KM8871 and KM8869, obtained in the item 6(2) of Example 1 which express various anti-GD3 CDR-grafted antibodies was suspended in GIT medium (manufactured by Nippon Pharmaceutical) containing 200 nM MTX to give a density of 1 to 2×10^5 cells/ml, and dispensed in 200 ml into 175 cm² flasks (manufactured by Greiner). The cells were cultured at 37°C for

5 to 7 days until they became confluent and then the culture supernatant was recovered. Each anti-GD3 CDR-grafted antibody was purified from the culture supernatant using Prosep-A (manufactured by Bioprocessing) column in accordance with the manufacture's instructions. About 3 mg of anti-GD3 CDR-grafted antibody KM8870 was obtained from 500 ml of the culture supernatant of KM8870, and about 25 mg of anti-GD3 CDR-grafted antibody KM8871 from 1,600 ml of the culture supernatant of KM8871 and about 65 mg of anti-GD3 CDR-grafted antibody KM8869 from 1,000 ml of the culture supernatant of KM8869. About 4 µg of each of the thus obtained anti-GD3 CDR-grafted antibodies was subjected to an electrophoresis in accordance with the known method (*Nature*, 227, 680 (1970)) to examine its molecular weight and purification degree. Results are shown in Fig. 13. As shown in Fig. 13, molecular weight of each purified anti-GD3 CDR-grafted antibody was about 150 kilodaltons (hereinafter referred to as "Kd") under non-reducing conditions, and two bands of about 50 Kd and about 25 Kd were found under reducing conditions. The molecular weights almost coincided with the molecular weights deduced from the cDNA nucleotide sequences of H chain and L chain of the antibody (H chain: about 49 Kd, L chain: about 24 Kd, whole molecule: about 146 Kd), and also coincided with the reports that the IgG type antibody has a molecular weight of about 150 Kd under non-reducing conditions and is degraded into H chains having a molecular weight of about 50 Kd and L chains having a molecular weight of about 25 Kd under reducing conditions due to cutting of the disulfide bond (hereinafter referred to as "S-S bond") in the molecule (*Antibodies: A Laboratory Manual, Monoclonal Antibodies: Principles and Practice*), so that it was confirmed that each anti-GD3 CDR-grafted antibody was expressed as an antibody molecule of a correct structure. Also, when N-terminal amino acid sequences of the H chain and L chain of each of the purified anti-GD3 CDR-grafted antibodies were analyzed by Edman degradation using a protein sequencer (470A, manufactured by Applied Biosystems), it was confirmed that they coincide with the H chain and L chain N-terminal amino acid sequences deduced from respective cDNA nucleotide sequences. (0167)

7. Evaluation of activity of anti-GD3 CDR-grafted antibody

(1) Reactivity of anti-GD3 CDR-grafted antibody with GD3 (ELISA)

Reactivity of purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 with GD3 (manufactured by DIA-IATRON) was measured by the ELISA shown in the item 2 of Example 1. Fig. 14 shows a result of the examination of the reactivity carried out by fixing the amount of GD3 to be adsorbed to each well of a plate for ELISA to 20 pmol/well and changing the concentration of anti-GD3 chimeric

antibody KM871 and anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 to be added. As shown in Fig. 14, among the anti-GD3 CDR-grafted antibodies, KM8871 showed the most highest binding activity which was about 1/2 of the anti-GD3 chimeric antibody KM871. Fig. 15 shows a result of the examination of the reactivity of a constant concentration (10 µg/ml) of anti-GD3 chimeric antibody KM871 and anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871, carried out by changing the amount of GD3 to be adsorbed to each well of a plate for ELISA use. As shown in Fig. 15, among the anti-GD3 CDR-grafted antibodies, KM8871 showed the most highest binding activity which was equivalent to the anti-GD3 chimeric antibody KM871. Fig. 16 shows a result of the examination of the reactivity of a constant concentration (10 µg/ml) of anti-GD3 chimeric antibody KM871 and anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871, carried out by changing the kind of ganglioside to be adsorbed to each well of a plate for ELISA use (adsorption amount: 20 pmol/well). A total of 11 kinds of ganglioside, GM1, N-acetylGM2 (manufactured by Boehringer Mannheim, hereinafter referred to as "AcGM2"), N-glycolylGM2 (hereinafter referred to as "GcGM2"), N-acetylGM3 (hereinafter referred to as "AcGM3"), N-glycolylGM3 (hereinafter referred to as "GcGM3"), GD1a and GD1b (manufactured by DIA-IATRON), GD2 and GD3 (manufactured by DIA-IATRON), GQ1b (manufactured by DIA-IATRON) and GT1b (manufactured by Funakoshi), were used. In this case, GM1 and GD1a were purified from bovine brain, and N-glycolylGM2 and N-glycolylGM3 from mouse liver, N-acetylGM3 from canine erythrocyte and GD2 from a human neuroblastoma culture cell line IMR32 (ATCC CCL 127), respectively in accordance with the known method (*J. Biol. Chem.*, 263, 10915 (1988)). As shown in Fig. 16, each of the anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 bound strongly to GD3 and weakly to GQ1b similar to the case of the anti-GD3 chimeric antibody KM871 but did not bind to other gangliosides. Thus, it was shown that they maintain the binding specificity of chimeric antibody KM871.

(0168)

(2) Reactivity of anti-GD3 CDR-grafted antibody with human cancer cell (immunofluorescent method)

Reactivity of purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 with human cancer cells was measured as follows. That is, 1×10^6 cells of each of human melanoma culture cell lines G-361 (ATCC CRL 1424) and SK-MEL-28 (ATCC HTB72) were suspended in PBS, put into microtubes and centrifuged (2,000 rpm for 2 minutes), and the thus washed cells were stirred after adding 50 µl of the anti-

GD3 chimeric antibody KM871 or the anti-GD3 CDR-grafted antibody, KM8869, KM8870 or KM8871 (a solution adjusted to 5 µg/ml with 1% BSA-PBS) and then allowed to react at 4°C for 1 hour. After the reaction, followed by centrifugation three times with PBS for washing, 20 µl of rabbit anti-human IgG (H + L) F(ab')₂ solution (manufactured by Wako Pure chemical Industries, used by diluting 30 times with 1% BSA-PBS) fluorescence-labeled with fluorescein isothiocyanate (hereinafter referred to as "FITC") was added thereto, followed by stirring, and then the reaction was carried out at 4°C for 1 hour. After the reaction, followed by centrifugation three times with PBS for washing, the cells were again suspended in PBS to carry out the analysis using a flow cytometer EPICS Elite (manufactured by COULTER). As a control, the analysis was carried out by the same procedure without adding antibodies. Results are shown in Fig. 17. As shown in Fig. 17, all of the anti-GD3 chimeric antibody KM871 and the anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 showed the reactivity with both cell lines. Among the anti-GD3 CDR-grafted antibodies, KM8871 showed the most strongest reactivity. The above results show that the anti-GD3 CDR-grafted antibodies are useful in the diagnosis, treatment and the like of GD3-positive human tumors including melanoma.

(0169)

(3) *In vitro* cytotoxic activity of anti-GD3 CDR-grafted antibodies (CDC activity)

In order to evaluate *in vitro* cytotoxic activity of the purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871, the CDC activity was measured by the following method.

a. Preparation of target cell suspension

Each of human melanoma culture cell lines, G-361 (ATCC CRL 1424) and SK-MEL-28 (ATCC HTB72), was cultured in RPMI1640-FBS(10) medium and adjusted to 5×10^6 cells, and the cells were isotope-labeled by adding 3.7 MBq equivalent of a radioactive substance $\text{Na}_2^{51}\text{CrO}_4$ and carrying out the reaction at 37°C for 1 hour. After the reaction, the cells were washed three times by their suspension in RPMI1640-FBS(10) medium and centrifugation, re-suspended in the medium and then incubated in ice at 4°C for 30 minutes to thereby spontaneously release the radioactive substance. After centrifugation, the precipitate was adjusted to 1×10^6 cells/ml by adding 5 ml of RPMI1640-FBS(10) medium and used as the target cell suspension.

(0170)

b. Preparation of complement solution

Sera of three healthy persons were mixed and used as the human complement source. At the time of its use, it was diluted to 15% vol/vol with RPMI1640-FBS(10) medium and used as the complement solution.

c. Measurement of CDC activity

To each well of a 96 well U bottom plate (manufactured by Falcon), 50 μ l of the target cell suspension prepared in "a" was added (5×10^4 cells/well), and then the anti-GD3 chimeric antibody KM871 or the anti-GD3 CDR-grafted antibody, KM8869, KM8870 or KM8871, was added to give a final concentration of 0.05 to 50 μ g/ml and allowed to react at room temperature for 30 minutes. After the reaction, the plate was centrifuged to remove the supernatant, and 150 μ l of the human complement solution prepared in "b" was added thereto and allowed to react at 37°C for 1 hour. After centrifugation, the amount of ^{51}Cr released in the supernatant was measured by a γ -counter. The amount of the spontaneously dissociated ^{51}Cr was calculated by carrying out the same procedure using the medium alone instead of the antibody solution and complement solution and measuring the amount of ^{51}Cr in the supernatant. Amount of the total dissociated ^{51}Cr was calculated by carrying out the same procedure by adding the medium alone instead of the antibody solution, and 5 N sodium hydroxide instead of the complement solution, and measuring the amount of ^{51}Cr in the supernatant. The CDC activity was calculated by the following equation.

(0171)

(Equation 1)

$$\text{CDC activity (\%)} = \frac{\begin{array}{l} \text{(amount of } ^{51}\text{Cr in sample supernatant)} \\ - \text{(amount of spontaneously released } ^{51}\text{Cr)} \end{array}}{\begin{array}{l} \text{(amount of total } ^{51}\text{Cr)} \\ - \text{(amount of spontaneously released } ^{51}\text{Cr)} \end{array}} \times 100$$

(0172)

Results are shown in Fig. 18. As shown in Fig. 18, it was found that all of the anti-GD3 chimeric antibody KM871 and the anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 show the CDC activity for both cell lines, particularly show remarkably strong cytotoxic activity for the human melanoma culture cell line G-361. Among the anti-GD3 CDR-grafted antibodies, KM8871 showed the most stronger cytotoxic activity which was about 1/3 of the anti-GD3 chimeric antibody KM871.

(0173)

(4) *In vitro* cytotoxic activity of anti-GD3 CDR-grafted antibody (ADCC activity)

In order to evaluate *in vitro* cytotoxic activity of the purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871, the ADCC activity was measured by the following method.

a. Preparation of target cell suspension

Each of human melanoma culture cell lines, G-361 (ATCC CRL 1424) and SK-MEL-28 (ATCC HTB72), was cultured in RPMI1640-FBS(10) medium and adjusted to 1×10^6 cells, and the cells were isotope-labeled by adding 1.85 MBq equivalent of a radioactive substance $\text{Na}_2^{51}\text{CrO}_4$ and carrying out the reaction at 37°C for 1 hour. After the reaction, the cells were washed three times by their suspension in RPMI1640-FBS(10) medium and centrifugation, re-suspended in the medium and then incubated in ice at 4°C for 30 minutes to thereby spontaneously release the radioactive substance. After centrifugation, the precipitate was adjusted to 2×10^5 cells/ml by adding 5 ml of RPMI1640-FBS(10) medium and used as the target cell suspension.

b. Preparation of effector cell suspension

Healthy human vein blood (50 ml) was collected, and 0.5 ml of heparin sodium (manufactured by Takeda Pharmaceutical) was gently added thereto. The resulting mixture was centrifuged ($1,500$ to $1,800 \times g$ for 30 minutes) using Polymorphprep (manufactured by Nycomed Pharma AS) in accordance with the manufacture's instructions to separate a mononuclear cell layer. The separated cells were three times centrifuged ($1,500$ to $1,800 \times g$ for 5 minutes) with RPMI1640-FBS(10) medium for washing and then re-suspended in the medium to give a density of 5×10^6 cells/ml to be used as the effector cell suspension.

(0174)

c. Measurement of ADCC activity

Into each well of a 96 well U bottom plate (manufactured by Falcon), 50 μl of the target cell suspension prepared in "a" was dispensed (1×10^4 cells/well). Next, 100 μl of the effector cell suspension prepared in "b" was added (5×10^5 cells/well, ratio of the effector cells to the target cells becomes 50:1). Thereafter, each of the anti-GD3 chimeric antibody KM871 and the anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 was added to give a final concentration of 0.05 to 50 $\mu\text{g}/\text{ml}$ and allowed to react at 37°C for 4 hours. After the reaction, the plate was centrifuged and the amount of ^{51}Cr in the supernatant was measured by a γ -counter. The amount of the spontaneously dissociated ^{51}Cr was calculated by carrying out the same procedure using the medium alone instead of the effector cell suspension and antibody solution and measuring the amount of ^{51}Cr in the supernatant. The amount of the total dissociated ^{51}Cr was calculated by carrying out the same procedure by adding the medium alone

instead of the antibody solution, and 5 N sodium hydroxide instead of the effector cell suspension, and measuring the amount of ^{51}Cr in the supernatant. The ADCC activity was calculated by the following equation.

(0175)

(Equation 2)

$$\text{ADCC activity (\%)} = \frac{\begin{array}{l} \text{(amount of } ^{51}\text{Cr in sample supernatant)} \\ - \text{(amount of spontaneously released } ^{51}\text{Cr)} \end{array}}{\begin{array}{l} \text{(amount of total } ^{51}\text{Cr)} \\ - \text{(amount of spontaneously released } ^{51}\text{Cr)} \end{array}} \times 100$$

(0176)

Results are shown in Fig. 19. As shown in Fig. 19, it was found that all of the anti-GD3 chimeric antibody KM871 and the anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871, show strong ADCC activity for both cell lines. Among the anti-GD3 chimeric antibody KM871 and the anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871, difference in the cytotoxic activity was not found.

(0177)

The above results show that the anti-GD3 CDR-grafted antibodies are useful in the diagnosis, treatment and the like of GD3-positive human tumors similar to the case of the anti-GD3 chimeric antibody KM871. In addition, it is expected that immunogenicity of the anti-GD3 CDR-grafted antibodies is reduced in human and the effect is further prolonged.

The transformed cell clone KM8871 capable of producing KM8871 which showed the most high activity among the anti-GD3 CDR-grafted antibodies has been deposited on July 22, 1999, as FERM BP-6790 in National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology (Higashi 1-1-3, Tsukuba, Ibaraki, Japan).

Example 2. Preparation of fusion protein of anti-GD3 CDR-grafted antibody with human cytokine:

As an example of a fusion protein of anti-GD3 CDR-grafted antibody with human cytokine, a fusion protein of the anti-GD3 CDR-grafted antibody KM8871 with human IL-2, KM8871-hIL-2, was prepared as follows and evaluation of the activity was carried out.

(0178)

1. Construction of cDNA encoding a fusion protein of hCy1 with hIL-2

(1) Construction of a plasmid pBSΔhCy1-IL-2 having a cDNA comprised of about 65 bases of 3'-terminal of hCy1 cDNA and full length cDNA of matured type hIL-2

A plasmid pBluescript SK(-) (3 µg, manufactured by Stratagene) was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed with 10 units of a restriction enzyme *Apal* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and further mixed with 10 units of a restriction enzyme *BamHI* (manufactured by Takara Shuzo) to carry out the reaction at 30°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an *Apal-BamHI* fragment of about 2.95 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0179)

Next, synthetic DNA fragments respectively having the nucleotide sequences represented by SEQ ID NO:49 and SEQ ID NO:50 were synthesized using an automatic DNA synthesizer (380A, manufactured by Applied Biosystems). A 0.3 µg of each of the thus obtained synthetic DNA fragments was added to 15 µl of sterile water and heated at 65°C for 5 minutes. After allowing the reaction solution to stand at room temperature for 30 minutes, this was mixed with 2 µl of a 10 × buffer solution (500 mM Tris-HCl (pH 7.6), 100 mM magnesium chloride, 50 mM DTT) and 2 µl of 10 mM ATP and further mixed with 10 units of T4 Polynucleotide Kinase (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 30 minutes to thereby phosphorylate the 5'-terminal.

(0180)

Next, 0.1 µg of the *Apal-BamHI* fragment derived from the plasmid pBluescript SK(-) and 0.05 µg of the phosphorylated synthetic DNA, both obtained in the above, were added to 10 µl in total volume of sterile water and ligated using DNA Ligation Kit Ver. 2 (manufactured by Takara Shuzo) in accordance with the manufacture's instructions. Using the thus obtained recombinant plasmid DNA solution, an *E. coli* line DH5α (manufactured by Stratagene) was transformed to obtain the plasmid pBSA-B shown in Fig. 20. The thus obtained plasmid (10 µg) was allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to an electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence, and as a result, it was confirmed that a plasmid into which the DNA of interest was cloned was obtained.

(0181)

Next, 3 µg of the thus obtained plasmid pBSA-B was added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed with 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 33 mM Tris-acetate (pH 7.9), 66 mM potassium acetate, 10 mM magnesium acetate, 0.5 mM DTT and 100 µg/ml BSA, and further mixed with 10 units of a restriction enzyme *SmaI* (manufactured by Takara Shuzo) to carry out the reaction at 30°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an *EcoRI-SmaI* fragment of about 3.00 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0182)

Next, the following PCR was carried out using a plasmid pILL4 containing full length cDNA of matured type hIL-2 (*Agric. Biol. Chem.*, 51, 1135 (1987)) as the template. The plasmid pILL4 (1 ng) was added to 100 µl of a reaction solution (1 × concentration Ex Taq buffer (manufactured by Takara Shuzo), 200 µM dNTPs, 1.0 µM rev1 primer (SEQ ID NO:51), 1.0 µM fw2 primer (SEQ ID NO:52) and 2.5 units of TaKaRa Ex Taq DNA polymerase (manufactured by Takara Shuzo)), and the solution was covered with 100 µl of mineral oil and set to a DNA thermal cycler (PJ480, manufactured by PERKIN ELMER) to carry out the reaction at 94°C for 3 minutes, subsequent 30 cycles of each cycle at 96°C for 30 seconds, 55°C for 1 minute and 72°C for 1 minute, and finally at 72°C for 7 minutes. The reaction solution was precipitated with ethanol, added to 30 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride and 1 mM DTT, and further mixed with 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 33 mM Tris-acetate (pH 7.9), 66 mM potassium acetate, 10 mM magnesium acetate, 0.5 mM DTT and 100 µg/ml BSA, and further mixed with 10 units of a restriction enzyme *SmaI* (manufactured by Takara Shuzo) to carry out the reaction at 30°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 1 µg of an *EcoRI-SmaI* fragment of about 0.41 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0183)

Next, 0.1 µg of the thus obtained *EcoRI-SmaI* fragment of the plasmid pBSA-B and 0.1 µg of the *EcoRI-SmaI* fragment of the PCR product of full length cDNA of matured type hIL-2 were added to 10 µl in total volume of sterile water and ligated using DNA Ligation Kit Ver. 2 (manufactured by Takara Shuzo) in accordance with the manufacture's instructions. Using the thus obtained recombinant plasmid DNA solution, an *E. coli* DH5α was transformed. Each plasmid DNA was prepared from 10 clones of the transformants, allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to an electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine nucleotide sequence of the inserted cDNA, and as a result, the plasmid pBSΔhCyl-IL-2 shown in Fig. 21 having the nucleotide sequence of interest was obtained.

(0184)

(2) Construction of a plasmid pBS Δ hCyl-IL-2 having a cDNA comprised of hCyl full length cDNA and full length cDNA of matured type hIL-2

First, 3 µg of the plasmid pBS Δ hCyl described in Japanese Published Unexamined Patent Application No. 257893/98 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed with 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and further mixed with 10 units of a restriction enzyme *EcoT22I* (manufactured by Takara Shuzo) to carry out the reaction at 30°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 1 µg of an *ApaI-EcoT22I* fragment of about 0.92 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0185)

Next, 3 µg of the plasmid pBSΔhCyl-IL-2 obtained in the item 1(1) of Example 2 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed with 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM sodium chloride, and further mixed with 10 units of a restriction enzyme *EcoT22I* (manufactured by Takara Shuzo) to carry out the reaction at 30°C for

1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an *ApaI-EcoT22I* fragment of about 3.40 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

5 (0186)

Next, 0.1 µg of the thus obtained *ApaI-EcoT22I* fragment of pBShCyl and 0.1 µg of the *ApaI-EcoT22I* of pBSΔhCyl-IL-2 were added to 10 µl in total volume of sterile water and ligated using DNA Ligation Kit Ver. 2 (manufactured by Takara Shuzo) in accordance with the manufacture's instructions. Using the thus obtained recombinant plasmid DNA solution, the *E. coli* DH5α was transformed to obtain the plasmid pBShCyl-IL-2 shown in Fig. 22. The thus obtained plasmid (10 µg) was allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to an electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine nucleotide sequence of the inserted cDNA, and as a result, it was confirmed that a plasmid having the nucleotide sequence of interest was obtained.

(0187)

2. Stable expression of KM8871-hIL-2 using animal cells

(1) Construction of stable expression vector for KM8871-hIL-2

A stable expression vector for KM8871-hIL-2 was constructed as follows using the vector pKANTEX641HLCDRm-69 obtained in the item 6(1) of Example 1 for stable expression of anti-GD3 CDR-grafted antibody KM8871 and the plasmid pBShCyl-IL-2 obtained in the item 1(2) of Example 2 having cDNA encoding the fusion protein of hCyl with hIL-2.

25 (0188)

The plasmid pKANTEX641HLCDRm-69 (3 µg) obtained in the item 6(1) of Example 1 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed with 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol and added to 10 µl of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and the solution was further mixed with 10 units of a restriction enzyme *BamHI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an *ApaI-BamHI* fragment of about

12.57 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0189)

Next, 3 µg of each of the plasmid pBShCyl-IL-2 obtained in the item 1(2) of Example 2 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and the solution was further mixed with 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol and added to 10 µl of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and the solution was further mixed with 10 units of a restriction enzyme *BamHI* (manufactured by Takara Shuzo) to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by an agarose gel electrophoresis to recover about 2 µg of an *ApaI*-*BamHI* fragment of about 1.45 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0190)

Next, 0.1 µg of the thus obtained *ApaI*-*BamHI* fragment of the plasmid pKANTEX641HLCDRIm-69 and 0.1 µg of the *ApaI*-*BamHI* fragment of the plasmid pBShCyl-IL-2 were added to 10 µl in total volume of sterile water and ligated using DNA Ligation Kit Ver. 2 (manufactured by Pharmacia) in accordance with the manufacture's instructions. Using the thus obtained recombinant plasmid DNA solution, an *E. coli* DH5α was transformed to obtain the KM8871-hIL-2 stable expression vector pKANTEX8871-hIL2 shown in Fig. 23. The thus obtained plasmid (10 µg) was allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to an electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence, and as a result, it was confirmed that a plasmid into which the DNA of interest was cloned was obtained.

(0191)

(2) Expression of KM8871-hIL-2 in animal cells

Using 4 µg of the KM8871-hIL-2 stable expression vector pKANTEX8871-hIL-2 obtained in the item 2(1) of Example 2, YB2/0 cell (ATCC CRL 1581) was transformed in accordance with the method described in the item 6(2) of Example 1, and selection was carried out finally with G418 (0.5 mg/ml) and MTX (200 nM) to obtain a transformed cell clone KM8871hIL2 showing an expression level of about 4 µg/10⁶ cells/24 hours. Also, the KM8871hIL2 has been deposited on July 22, 1999, as

FERM BP-6791 in National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology (Higashi 1-1-3, Tsukuba, Ibaraki, Japan).

(0192)

(3) Purification of KM8871-hIL-2 from culture supernatant

5 In accordance with the method described in the item 6(3) of Example 1, the transformed cell clone KM8871hIL2 obtained in the item 2(2) of Example 2 capable of expressing KM8871-hIL-2 was cultured to obtain about 10.0 mg of purified KM8871-hIL-2 from about 3 L of the culture supernatant. A result of SDS-PAGE of the purified KM8871-hIL-2 is shown in Fig. 24. As shown in Fig. 24, molecular weight of the purified KM8871-hIL-2 was about 180 Kd under non-reducing conditions, and two bands of about 65 Kd and about 25 Kd were found under reducing conditions. These molecular weights almost coincided with the molecular weights deduced from the cDNA nucleotide sequences of H chain of KM8871-hIL-2 and hIL-2 and L chain (H chain and hIL-2: about 64 Kd, L chain: about 24 Kd, whole molecule: about 176 Kd), and it was confirmed that the structure as antibody molecule was maintained after fusion of hIL-2.

(0193)

3. Evaluation of activity of KM8871-hIL-2

(1) Reactivity of KM8871-hIL-2 with GD3 (ELISA method)

20 Reactivity of the purified KM8871-hIL-2 with GD3 (manufactured by DIA-IATRON) was measured in accordance with the method described in the item 2 of Example 1. In this case, a peroxidase-labeled goat anti-human IgG (H & L) antibody (manufactured by American Qualex, used by diluting 3,000 times with 1% BSA-PBS) was used as the secondary antibody solution. Fig. 25 shows a result of the examination of the reactivity carried out by fixing the amount of GD3 to be adsorbed to each well of a plate for ELISA use to 20 pmol/well and changing the concentration of the anti-GD3 CDR-grafted antibody KM8871 and KM8871-hIL-2 to be added. As shown in Fig. 25, it was found that KM8871-hIL-2 has a GD3 binding activity similar to or higher than that of the anti-GD3 CDR-grafted antibody KM8871. Fig. 26 shows a result of the examination of the reactivity of a constant concentration (10 µg/ml) of anti-GD3 CDR-grafted antibody KM8871 and KM8871-hIL-2, carried out by changing the kinds of ganglioside to be adsorbed to each well of a plate for ELISA use (adsorption amount: 20 pmol/well). As shown in Fig. 26, it was found that the KM8871-hIL-2 strongly binds to GD3 similar to the case of the anti-GD3 CDR-grafted antibody KM8871, and it was observed that KM8871-hIL-2 also strongly binds to GD2. As the cause of the cross-reactivity of KM8871-hIL-2 with GD2, a possibility of the

binding of hIL-2 with GD2, as well as a possibility of the change in binding specificity caused by a change in the three-dimensional structure of antibody V region due to fusion of hIL2, a possibility of the change in the reactivity of secondary antibody caused by a change in the three-dimensional structure of antibody C region due to fusion of hIL2, and the like can be considered.

(0194)

(2) Reactivity of KM8871-hIL-2 with cell surface GD3 (an immunofluorescent method)

Reactivity of the purified KM8871-hIL-2 with human cancer cells was measured in accordance with the method described in the item 7(2) of Example 1. Results are shown in Fig. 27. As shown in Fig. 27, KM8871-hIL-2 showed a strong reactivity with the human melanoma culture cell line G361, which was almost identical to that of the anti-GD3 CDR-grafted antibody KM8871. This result shows that the KM8871-hIL-2 is useful in the treatment and the like of GD3-positive human tumors including melanoma.

(0195)

(3) Evaluation of hIL-2 activity of KM8871-hIL-2

Activity of the purified KM8871-hIL-2 as hIL-2 was measured in accordance with the following method. A mouse T cell line CTLL-2 (ATCC TIB214) of which growth shows dependency on hIL-2 concentration was suspended in RPMI1640-FBS(10) medium at a density of 2×10^5 cells/ml and dispensed in 50 μ l/well into a 96 well microtiter plate (manufactured by Sumitomo Bakelite). A solution (50 μ l) prepared by diluting hIL-2 (manufactured by R & D SYSTEMS) or purified KM8871-hIL-2 to various concentrations with RPMI1640-FBS(10) medium was added to each well and cultured at 37°C for 30 hours in a 5% CO₂ incubator. After the culturing the number of intact cells was counted using Cell Counting Kit (manufactured by Dojindo Laboratories) in accordance with the manufacture's instructions. Results are shown in Fig. 28. As shown in Fig. 28, KM8871-hIL-2 showed a biological activity to support CTLL-2 growth similar to that of hIL-2. These results show that the activity of KM8871-hIL-2 as hIL-2 is maintained after its fusion with the anti-GD3 CDR-grafted antibody KM8871.

(0196)

(4) Activation of human effector cell and enhancement of cytotoxic activity by KM8871-hIL-2

In order to evaluate activation of human effector cells and accompanying enhancement of cytotoxic activity by the hIL-2 moiety of KM8871-hIL-2 *in vitro*, the cytotoxic activity was measured by the following method.

a. Preparation of target cell suspension

In accordance with the method described in "a" of the item 7(4) of Example 1, a human melanoma culture cell line G-361 was adjusted to a density of 2×10^5 cells/ml and used as the target cell suspension.

5 b. Preparation of human effector cell suspension

In accordance with the method described in "b" of the item 7(4) of Example 1, mononuclear cells were separated from venous blood of healthy person and re-suspended to a density of 5×10^6 cells/ml to be used as the effector cell suspension.

(0197)

10 c. Activation of human effector cells

The effector cell suspension (100 μ l) prepared in "b" was dispensed into each well of a 96 well U bottom plate (manufactured by Falcon) (5×10^4 cells/well). Furthermore, 50 μ l of the anti-GD3 CDR-grafted antibody KM8871 or KM8871-hIL-2 was added thereto to give a final concentration of 11.1 nM and allowed to react at 37°C for 24 hours in a 5% CO₂ incubator.

d. Measurement of cytotoxic activity

The target cell suspension (50 μ l) prepared in "a" was added to each well of the plate prepared in "c" (1×10^4 cells/well). In this case, the ratio of the effector cells to the target cells becomes 5:1. After the reaction at 37°C for 4 hours, the plate was centrifuged and the amount of ⁵¹Cr in the supernatant was measured by a γ -counter. The amount of the spontaneously dissociated ⁵¹Cr was calculated by carrying out the same procedure using the medium alone instead of the effector cell suspension and antibody solution and measuring the amount of ⁵¹Cr in the supernatant. The amount of the total dissociated ⁵¹Cr was calculated by carrying out the same procedure by adding the medium alone instead of the antibody solution, and 1 N hydrochloric acid instead of the effector cell suspension, and measuring the amount of ⁵¹Cr in the supernatant. The cytotoxic activity was calculated by the following equation.

(0198)

(Equation 3)

30

$$\text{ADCC activity (\%)} = \frac{\begin{array}{l} \text{(amount of } ^{51}\text{Cr in sample supernatant)} \\ - \text{(amount of spontaneously released } ^{51}\text{Cr)} \end{array}}{\begin{array}{l} \text{(amount of total } ^{51}\text{Cr)} \\ - \text{(amount of spontaneously released } ^{51}\text{Cr)} \end{array}} \times 100$$

35 (0199)

Results are shown in Fig. 29. As shown in Fig. 29, it was found that the cytotoxic activity induced by KM8871-hIL-2 was stronger than that induced by the anti-GD3 CDR-grafted antibody KM8871. This result shows that human effector cells can

be activated and the cytotoxic activity can be enhanced by using KM8871-hIL-2. Thus, it was indicated that similar effects can be expected in its clinical application to human.

(0200)

Example 3. Production of fusion protein of anti-GD3 chimeric antibody with human cytokine:

As an example of fusion protein of anti-GD3 chimeric antibody with human cytokine, a fusion protein of the anti-GD3 chimeric antibody KM871 with human IL-2, KM871-hIL-2, was prepared as follows to carry out evaluation of the activity.

1. Stable expression of KM871-hIL-2 using animal cells

(1) Construction of stable expression vector for KM871-hIL-2

A stable expression vector for KM871-hIL-2 was constructed as follows using the vector pKANTEX641 for stable expression of anti-GD3 chimeric antibody KM871 (Japanese Published Unexamined Patent Application No. 304989/93) and the plasmid pBShCyl-IL-2 obtained in the item 1(2) of Example 2 having cDNA encoding the fusion protein of hCyl with hIL-2.

(0201)

The plasmid pKANTEX641 (3 µg) obtained in the item 6(1) of Example 1 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol and added to 10 µl of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and 10 units of a restriction enzyme *BamHI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 30°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 2 µg of an *ApaI*-*BamHI* fragment of about 12.57 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0202)

Next, 3 µg of each of the plasmid pBShCyl-IL-2 obtained in the item 1(2) of Example 2 was added to 10 µl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 10 mM magnesium chloride and 1 mM DTT, and 10 units of a restriction enzyme *ApaI* (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol and added to 10 µl of a buffer comprising 20 mM Tris-HCl (pH 8.5), 10 mM magnesium chloride, 1 mM DTT and 100 mM potassium chloride, and 10 units of a restriction enzyme *BamHI*

(manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was fractionated by agarose gel electrophoresis to recover about 2 µg of an *ApaI-BamHI* fragment of about 1.45 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0203)

Next, 0.1 µg of the thus obtained *ApaI-BamHI* fragment of the plasmid pKANTEX641 and 0.1 µg of the *ApaI-BamHI* fragment of the plasmid pBShCyl-IL-2 were added to 10 µl in total volume of sterile water and ligated using DNA Ligation Kit Ver. 2 (manufactured by Pharmacia) in accordance with the manufacture's instructions. Using the thus obtained recombinant plasmid DNA solution, an *E. coli* DH5α was transformed to obtain the KM871-hIL-2 stable expression vector pKANTEX871-hIL2 shown in Fig. 30. The thus obtained plasmid (10 µg) was allowed to react in accordance with the manufacture's instructions attached to AutoRead Sequencing Kit (manufactured by Pharmacia) and then subjected to electrophoresis using A.L.F. DNA Sequencer (manufactured by Pharmacia) to determine the nucleotide sequence, and as a result, it was confirmed that a plasmid into which the DNA of interest was cloned was obtained.

(0204)

(2) Expression of KM871-hIL-2 in animal cells

Using 4 µg of the KM871-hIL-2 stable expression vector pKANTEX871-hIL-2 obtained in the item 1(1) of Example 3, YB2/0 cell (ATCC CRL 1581) was transformed in accordance with the method described in the item 6(2) of Example 1, and selection was carried out finally with G418 (0.5 mg/ml) and MTX (200 nM) to obtain a transformed cell clone KM871hIL2 showing an expression level of about 4 µg/10⁶ cells/24 hours. Also, the KM871hIL2 has been deposited on October 19, 1999, as FERM BP-6918 in National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology (Higashi 1-1-3, Tsukuba, Ibaraki, Japan).

(0205)

(3) Purification of KM871-hIL-2 from culture supernatant

In accordance with the method described in the item 6(3) of Example 1, the transformed cell clone KM871hIL2 obtained in the item 1(2) of Example 3 which expresses KM871-hIL-2 was cultured to obtain about 10.0 mg of purified KM871-hIL-2 from about 3 L of the culture supernatant. A result of SDS-PAGE of the purified KM871-hIL-2 is shown in Fig. 31. As shown in Fig. 31, molecular weight of the purified KM871-hIL-2 was about 180 Kd under non-reducing conditions, and two bands

of about 65 Kd and about 25 Kd were found under reducing conditions. These molecular weights almost coincided with the molecular weights deduced from the cDNA nucleotide sequences of H chain of KM871-hIL-2 and hIL-2 and L chain (H chain and hIL-2: about 64 Kd, L chain: about 24 Kd, whole molecule: about 176 Kd), and it was confirmed that the structure as antibody molecule was maintained after fusion of hIL-2.

(0206)

2. *In vitro* evaluation of KM871-hIL-2

(1) Reactivity of KM871-hIL-2 with GD3 (ELISA method)

Reactivity of the purified KM871-hIL-2 with GD3 (manufactured by DIA-IATRON) was measured in accordance with the method described in the item 2 of Example 1. In this case, a peroxidase-labeled goat anti-human IgG (H & L) antibody (manufactured by American Qualex, used by diluting 3,000 times with 1% BSA-PBS) was used as the secondary antibody solution. Fig. 32 shows results of the examination of the reactivity carried out by fixing the amount of GD3 to be adsorbed to each well of a plate for ELISA to 20 pmol/well and changing the concentration of the anti-GD3 chimeric antibody KM871 and KM871-hIL-2 to be added. As shown in Fig. 32, it was found that KM871-hIL-2 has a GD3 binding activity similar to or higher than that of the anti-GD3 chimeric antibody KM871. Fig. 33 shows results of the examination of the reactivity of a constant concentration (6.7 nM) of anti-GD3 chimeric antibody KM871 and KM871-hIL-2, carried out by changing the kinds of ganglioside to be adsorbed to each well of a plate for ELISA use (adsorption amount: 20 pmol/well). As shown in Fig. 33, it was found that the KM871-hIL-2 strongly binds to GD3 similar to the anti-GD3 chimeric antibody KM871, and it was confirmed that the addition of hIL-2 does not have large influence on the antigen specificity of KM871.

(0207)

(2) Reactivity of KM871-hIL-2 with cell surface GD3 (immunofluorescent method)

Reactivity of the purified KM871-hIL-2 with human cancer cells was measured in accordance with the method described in the item 7(2) of Example 1. Results are shown in Fig. 34. As shown in Fig. 34, KM871-hIL-2 showed a strong reactivity with the human melanoma culture cell line G361 and SK-MEL-28, which was almost identical to that of the anti-GD3 chimeric antibody KM871. The results show that the KM871-hIL-2 is useful in the treatment and the like of GD3-positive human tumors including melanoma.

(0208)

(3) Evaluation of hIL-2 activity of KM871-hIL-2

Activity of the purified KM871-hIL-2 as hIL-2 was measured in accordance with the item 3(2) of Example 2, and results are shown in Fig. 35. As shown in Fig. 35, KM871-hIL-2 showed a biological activity to support CTLL-2 growth similar to that of hIL-2. The results show that the activity of KM871-hIL-2 as hIL-2 is maintained after its fusion with the anti-GD3 chimeric antibody KM871.

(0209)

(4) Activation of human lymphocyte by KM871-hIL-2

In order to evaluate activation of human lymphocyte and accompanying enhancement of cytotoxic activity by the hIL-2 moiety of KM871-hIL-2 *in vitro*, the cytotoxic activity was measured by the following method in which the method described in the item 7(4) of Example 1 was partially changed.

a. Preparation of target cell suspension

In accordance with the method described in "a" of the item 7(4) of Example 1, a human melanoma culture cell line G-361 was adjusted to give a density of 2×10^5 cells/ml and used as the target cell suspension.

b. Preparation of effector cell suspension

In accordance with the method described in "b" of the item 7(4) of Example 1, mononuclear cells were separated from venous blood of healthy person and re-suspended to give a density of 1×10^7 cells/ml to be used as the effector cell suspension.

c. Activation of lymphocyte

The effector cell suspension (50 μ l) prepared in "b" was dispensed into each well of a 96 well U bottom plate (manufactured by Falcon) (5×10^4 cells/well). Furthermore, 50 μ l of RPMI1640-FBS(10) medium to which the anti-GD3 chimeric antibody KM871 or KM871-hIL-2 had been added was added thereto to give a final concentration of 1 nM (molecular weight of KM871 was calculated as 150 Kd, and molecular weight of KM871-hIL-2 as 180 Kd) and incubated at 37°C for 72 hours in a 5% CO₂ incubator.

(0210)

d. Measurement of cytotoxic activity

The target cell suspension (50 μ l) prepared in "a" was added to each well of the plate prepared in "c" (1×10^4 cells/well). In this case, the ratio of the effector cells to the target cells becomes 5:1. Furthermore, 50 μ l of RPMI1640-FBS(10) medium to which the anti-GD3 chimeric antibody KM871 or KM871-hIL-2 had been added was added thereto to give a final concentration of 1 nM to carry out the reaction at 37°C for 4 hours, and then the plate was centrifuged and the amount of ⁵¹Cr in the supernatant was measured by a γ -counter. The amount of the spontaneously dissociated ⁵¹Cr was

calculated by carrying out the same procedure using the medium alone instead of the effector cell suspension and antibody solution and measuring the amount of ^{51}Cr in the supernatant. The amount of the total dissociated ^{51}Cr was calculated by carrying out the same procedure by adding the medium alone instead of the antibody solution, and 1 N hydrochloric acid instead of the effector cell suspension, and measuring the amount of ^{51}Cr in the supernatant. The cytotoxic activity was calculated by the method described in "b" of the item 7(4) of Example 1, and results are shown in Fig. 36. As shown in Fig. 36, it was found that a cytotoxic activity induced by KM871-hIL-2 was stronger than that induced by KM871. Based on the results, it is expected that the KM871-hIL-2 will also enhance therapeutic effects of KM871 in the clinical application of GD3-positive human tumors by the activation of human lymphocyte via the hIL-2 moiety.

(0211)

3. *In vivo* evaluation of KM871-hIL-2

(1) Production of B16 cell transfectants introduced with GD3 synthase gene for mouse syngenic model

a. Construction of GD3 synthase gene stable expression vector

A vector pAMo-GD3 (3 μg ; WO 94/23020) for expressing GD3 synthase gene in animal cell was added to 10 μl of a buffer comprising 10 mM Tris-HCl (pH 7.5), 50 mM sodium chloride, 10 mM magnesium chloride and 1 mM DTT, and 10 units of a restriction enzyme *Hind*III (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol and added to 10 μl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride, 1 mM DTT, 100 $\mu\text{g}/\text{ml}$ BSA and 0.01% Triton X-100, and 10 units of a restriction enzyme *Not*I (manufactured by Takara Shuzo) were further added thereto to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, and the 5' protruding end formed by the restriction enzyme digestion was changed to blunt end using DNA Blunting Kit (manufactured by Takara Shuzo). The reaction solution was fractionated by agarose gel electrophoresis to recover about 2 μg of a blunt-ended *Hind*III-*Not*I fragment of about 2.1 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

Next, 3 μg of the anti-GD3 chimeric antibody expression vector pKANTEX641 was added to 10 μl of a buffer comprising 50 mM Tris-HCl (pH 7.5), 100 mM sodium chloride, 10 mM magnesium chloride and 1 mM DTT, and 10 units of a restriction enzyme *Eco*RI (manufactured by Takara Shuzo) were further added thereto

to carry out the reaction at 37°C for 1 hour. The reaction solution was precipitated with ethanol, and the 5' protruding end formed by the restriction enzyme digestion was changed to blunt end using DNA Blunting Kit (manufactured by Takara Shuzo). The reaction solution was precipitated with ethanol, added to 10 µl of a buffer comprising 50 mM Tris-HCl (pH 7.5) and 10 mM magnesium chloride, and the solution was further mixed with 10 units of a 5'-terminal dephosphorylation enzyme, calf intestine-derived alkaline phosphatase (manufactured by Takara Shuzo), and allowed to react at 37°C for 1 hour for dephosphorylation of the 5' end. The reaction solution was subjected to phenol treatment and then fractionated by agarose gel electrophoresis to recover about 2 µg of a blunt end-dephosphorylated *EcoRI-NruI* fragment of about 9.4 kb using QIAquick Gel Extraction Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(0212)

Next, 0.1 µg of the *EcoRI-NruI* fragment derived from the plasmid pKANTEX641 and 0.1 µg of the *HindIII-NofI* fragment derived from the plasmid pAMoGD3, both obtained in the above, were added to 10 µl in total volume of sterile water and ligated using DNA Ligation Kit Ver. 2 (manufactured by Takara Shuzo) in accordance with the manufacture's instructions. Using the thus obtained recombinant plasmid DNA solution, an *E. coli* DH5α was transformed to obtain the GD3 synthase-stable expression vector pKANTEXGD3 shown in Fig. 37.

(0213)

b. Introduction of GD3 synthase-expression vector into B16 cells

Using 4 µg of the GD3 synthase gene-stable expression vector pKANTEXGD3 obtained in "a" of the item 3(1) of Example 3, B16-F10 cell (ATCC CRL 1581) was transformed in accordance with the method described in the item 6(2) of Example 1, and selection was carried out finally by G418 (0.5 mg/ml) to obtain transfectant clones B16-8-3 and B16-29-10.

(0214)

c. Analysis of GD3 expression level of GD3-expressing line by immunofluorescent method

Each of 1×10^6 cells of the GD3 synthase gene transfectants B16-8-3 and B16-29-10 obtained in "b" of the item 3(1) of Example 3 and the parent line B16-F10 cell were suspended in PBS, put into microtubes and centrifuged (2,000 rpm for 2 minutes), and the thus washed cells were stirred after adding 50 µl of the anti-GD3 chimeric antibody KM871 (a solution prepared by adjusting to 5 µg/ml with 1% BSA-PBS, negative control was 1% BSA-PBS alone) and then allowed to react at 4°C for 1

hour. After the reaction, followed by centrifugation three times with PBS for washing, the cells were mixed with 100 μ l of biotin-labeled goat anti-human IgG (H + L) solution (manufactured by VECTOR, used by diluting 400 times with 1% BSA-PBS), stirred and then allowed to react at 4°C for 1 hour. After centrifugation three times with PBS for washing, 100 μ l of a streptavidin-labeled Cy5 pigment solution (Streptavidin-RED670 manufactured by GIBCO, used by diluting 100 times with 1% BSA-PBS) was added and stirred and then allowed to react at 4°C for 1 hour. After the reaction, followed by centrifugation three times with PBS for washing, the cells were again suspended in PBS to carry out the analysis using a flow cytometer EPICS Elite (manufactured by COULTER). Results are shown in Fig. 38. As shown in Fig. 38, the anti-GD3 chimeric antibody KM871 reacted only with B16-8-3 and B16-29-10 but did not show reactivity with the parent line B16-F10 cells. The above results show that both of the transfectants B16-8-3 and B16-29-10 express GD3 on the surface and can be used as a model for measuring *in vivo* antitumor effects of KM871-hIL-2 by transplanting them into C57 BL/6 mouse as the origin of B16 cells.

(0215)

(2) Measurement of *in vivo* antitumor effects of KM871-hIL-2

a. Evaluation with mice transplanted with GD3-expressing transformant B16-29-10

The GD3-expressing B16 cell B16-29-10 obtained in "b" of the item 3(1) of Example 3 was suspended in PBS to give a density of 2.5×10^6 cells/ml, and 100 μ l of the suspension was injected into a C57 BL/6 Cr slc mouse (female, 8 week-old, available from Japan SLC) through its tail vein. Also, starting on the day of tumor transplantation, 50 or 200 μ g/day of KM871 or 20 or 50 μ g/day of KM871-hIL-2 was administered intravenously once a day continuously for 5 days. The lungs were excised on the 15th day after the tumor transplantation, and visible black metastatic foci on the surface were counted, and results are shown in Table 3 and Fig. 39.

(0216)

(Table 3)

Table 3

Group		The number of metastatic foci on the lung surface	Average value
Non-administered group		147/136/106/72/70	106.2
KM871-hIL-2	20 μ g	20/15/11/2/2/1	8.5
KM871-hIL-2	50 μ g	15/12/5/2/0/0	5.7
KM871	50 μ g	185/114/86/77	115.5
KM871	200 μ g	113/87/86/86/77/55	84

(0217)

As shown in Table 3 and Fig. 39, KM871-hIL-2 showed clearly more potent anti-metastatic effect than that of KM871 in the model mouse of the clone B16-29-10.

b. Evaluation with mice transplanted with GD3-expressing transformant B16-8-3

Using the GD3-expressing B16 cell B16-8-3 obtained in "b" of the item 3(1) of Example 3, anti-metastatic effect of KM871-hIL-2 was measured in accordance with the method described in "a" of the item 3(2) of Example 3. The dose was 40 or 100 µg/day for both KM871 and KM871-hIL-2. Results are shown in Table 4 and Fig. 40.

(0218)

(Table 4)

Table 4

Group	The number of metastatic foci on the lung surface	Average value
Non-administered group	170/128/113/66/42/16/0	76.4
KM871-hIL-2 40 µg	1/0/0	0.3
KM871-hIL-2 100 µg	3/0/0	1
KM871 40 µg	128/70/30	76
KM871 100 µg	26/20/17	21

(0219)

As shown in Table 4 and Fig. 40, KM871-hIL-2 showed more potent anti-metastatic effect than that of KM871 in the model mouse of the clone B16-8-3.

(0220)

c. Comparison with combined administration of KM871 and hIL-2

The GD3-expressing B16 cells B16-29-10 obtained in "b" of the item 3(1) of Example 3 were suspended in PBS to give a density of 2.5×10^6 cells/ml, and 100 µl of the suspension was injected into a C57 BL/6 Cr slc mouse (female, 8 week-old, available from Japan SLC) through its tail vein. Also, starting on the day of tumor transplantation, 6 µg/day (0.0333 nmol) or 18 µg/day (0.1 nmol) of KM871-hIL-2 was administered intravenously once a day continuously for 5 days. Using other groups, a mixed solution of KM871 and hIL-2 (manufactured by Peprtech) corresponding to the amount of the administered KM871-hIL-2 (5 µg (0.0333 nmol) of KM871 and 1 µg (0.0667 nmol) of hIL-2, or 15 µg (0.1 nmol) of KM871 and 3 µg (0.2 nmol) of hIL-2) was administered through the tail vein once a day continuously for 5 days, starting on the day of tumor transplantation. The lungs were excised on the 15th day after the tumor transplantation, and visible black metastatic foci on the surface were counted, and results are shown in Table 5 and Fig. 41.

(0219)

(Table 5)

Table 5

Group		The number of metastatic foci on the lung surface	Average value
Non-administered group		163/132/97/95/88/57	105.3
KM871-hIL-2	6 μ g	26/26/17/12/8	17.8
KM871-hIL-2	18 μ g	43/34/28/13/2	24
KM871 + hIL-2	5 μ g + 1 μ g	215/117/97/85/56	114
KM871 + hIL-2	15 μ g + 3 μ g	92/61/48/46/42	57.8

(0222)

As shown in Table 5 and Fig. 41, KM871-hIL-2 showed more potent anti-metastatic effect than that of the combined administration of the equivalent amount of KM871 and hIL-2 in the model mouse of the clone B16-29-10. The results show that the *in vivo* antitumor effect shown by KM871-hIL-2 is not a simple additive effect of KM871 and hIL-2 but a result of the efficient induction of antitumor immunity due to activation of peripheral lymphocytes by the hIL-2 moiety of KM871-hIL-2 accumulated in the tumor, thus showing a possibility that KM871-hIL-2 will become a therapeutic agent having higher therapeutic effect than that of the conventional administration of hIL-2 alone or its combination use with an antibody and also having reduced side effects derived from hIL-2.

(0223)

(Effects of the invention)

According to the present invention, a human CDR-grafted antibody which specifically reacts with GD3 or the antibody fragment thereof and derivatives of an antibody against ganglioside GD3 or the antibody fragment thereof are provided.

(0224)

(Sequence listing)

SEQUENCE LISTING

<110> KYOWA HAKKO KOGYO CO., LTD

<120> Humanized anti-GD3 antibody and it's cytokine conjugate

<130>

<140>

<141>

<160> 56

<170> PatentIn Ver. 2.0

<210> 1

<211> 138

<212> PRT

<213> Mus musculus

<400> 1

5 Met Glu Phe Gly Leu Ser Trp Leu Phe Leu Val Leu Val Phe Lys Gly
-19 -15 -10 -5

Val Gln Cys Glu Val Thr Leu Val Glu Ser Gly Gly Asp Phe Val Lys
-1 1 5 10

10 Pro Gly Gly Ser Leu Lys Val Ser Cys Ala Ala Ser Gly Phe Ala Phe
15 20 25

15 Ser His Tyr Ala Met Ser Trp Val Arg Gln Thr Pro Ala Lys Arg Leu
30 35 40 45

Glu Trp Val Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser
50 55 60

20 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
65 70 75

Thr Leu Tyr Leu Gln Met Arg Ser Leu Arg Ser Glu Asp Ser Ala Met
80 85 90

25 Tyr Phe Cys Thr Arg Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser Trp
95 100 105

30 Gly Gln Gly Thr Thr Leu Thr Val Ser Ser
110 115

(0225)

35 <210> 2

<211> 128

<212> PRT

<213> Mus musculus

<400> 2

40 Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
-20 -15 -10 -5

Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Thr Ala Ser Ser Leu Pro
-1 1 5 10

45 Ala Ser Leu Gly Asp Arg Val Thr Ile Ser Cys Ser Ala Ser Gln Asp
15 20 25

50 Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val
30 35 40

Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
45 50 55 60

55 Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser

	65	70	75
	Asn Leu Glu Pro Glu Asp Ile Ala Thr Tyr Phe Cys His Gln Tyr Ser		
	80	85	90
5	Lys Leu Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg		
	95	100	105
10	(0226)		
	<210> 3		
	<211> 5		
	<212> PRT		
15	<213> Mus musculus		
	<400> 3		
	His Tyr Ala Met Ser		
	1		
20			
	<210> 4		
	<211> 17		
	<212> PRT		
	<213> Mus musculus		
25	<400> 4		
	Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val Lys		
	1 5 10 15		
	Gly		
30			
	<210> 5		
	<211> 10		
	<212> PRT		
35	<213> Mus musculus		
	<400> 5		
	Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser		
	1 5		
40			
	<210> 6		
	<211> 11		
	<212> PRT		
	<213> Mus musculus		
45	<400> 6		
	Ser Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn		
	1 5 10		
50			
	<210> 7		
	<211> 7		
	<212> PRT		
	<213> Mus musculus		
	<400> 7		
55	Tyr Ser Ser Asn Leu His Ser		

1 5

5 <210> 8
 <211> 9
 <212> PRT
 <213> Mus musculus
 <400> 8
 His Gln Tyr Ser Lys Leu Pro Trp Thr
 10 1 5

(0227)

15 <210> 9
 <211> 119
 <212> PRT
 <213> Artificial Sequence
 <400> 9
 20 Glu Val Gln Leu Val Glu Ser Gly Gly Asp Phe Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Val Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser His Tyr
 20 25 30
 25 Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 30 Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 35 Leu Gln Met Arg Ser Leu Arg Ala Glu Asp Ser Ala Val Tyr Phe Cys
 85 90 95
 Thr Arg Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser Trp Gly Gln Gly
 100 105 110
 40 Thr Leu Leu Thr Val Ser Ser
 115

45 (0228)

50 <210> 10
 <211> 108
 <212> PRT
 <213> Artificial Sequence
 <400> 10
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 55 Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr

20 25 30
 Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45
 5 Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 10 Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys His Gln Tyr Ser Lys Leu Pro Trp
 85 90 95
 15 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
 100 105

 (0229)
 20 <210> 11
 <211> 95
 <212> DNA
 <213> Artificial Sequence
 25 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 11
 caggaaacag ctatgacgcg gccgccacca tggagtttgg gctcagctgg ctttttcttg 60
 30 tccttgtttt caaagggtgtt cagtgtgagg tgcag 95

 <210> 12
 <211> 97
 35 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 12
 40 aaagcgaatc cagaggctgc acaggagact ctcagagacc ccccggtg tacaaagtct 60
 ccccgagact ccaccagctg cacctcacac tgaacac 97

 45 <210> 13
 <211> 98
 <212> DNA
 <213> Artificial Sequence
 <220>
 50 <223> Description of Artificial Sequence: synthetic DNA
 <400> 13
 gcagcctctg gattcgcttt cagtcattat gccatgtctt gggtccgcca ggctccaggg 60
 aagggsgctgg agtgggtggc ttatattagt agtgggtg 98
 55

<210> 14
 <211> 97
 <212> DNA
 5 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 14
 10 gatacagcgt gttcttggag ttatctctgg agatggtgaa tctgcccttt acactgtctg 60
 aatagtaggt gccactacca ccactactaa tataagc 97

(0230)
 15 <210> 15
 <211> 98
 <212> DNA
 <213> Artificial Sequence
 20 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 15
 ctccaagaac acgctgtatc tgcaaatgcg cagcctgaga gctgaggact cggctgtgta 60
 25 tttctgtaca agagtttaac tgggaaccta ctactttg 98

<210> 16
 <211> 92
 30 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 16
 35 gttttccag tcacgacggg cccttgggtg aggctgagga gacggtgagc agggttccct 60
 ggccccagga gtcaaagtag taggttccca gt 92

40 <210> 17
 <211> 94
 <212> DNA
 <213> Artificial Sequence
 <220>
 45 <223> Description of Artificial Sequence: synthetic DNA
 <400> 17
 caggaaacag ctatgacgaa ttccaccatg atgtcctctg ctacgttcct tggctcctg 60
 ttgctctgtt ttaaggtac cagatgtgac atcc 94
 50

<210> 18
 <211> 87
 <212> DNA
 55 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 18
 5 gcactacaag tgatggtagac tctgtctcct acagatgcag acagggagga tggagactgg 60
 gtcactctgga tgcacatct ggtacct 87

(0231)
 10 <210> 19
 <211> 89
 <212> DNA
 <213> Artificial Sequence
 15 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 19
 gtcacatca cttgtagtgc aagtcaggac attagtaatt atttaaactg gtatcagcag 60
 20 aaaccaggga aagcccctaa gctcctgat 89

<210> 20
 <211> 89
 25 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 20
 30 taatctgtcc cagatccacc gccgctgaac cttgatggga cccccgagtg taaatttgat 60
 gagtaaaaga tcaggagctt aggggcttt 89

35 <210> 21
 <211> 92
 <212> DNA
 <213> Artificial Sequence
 <220>
 40 <223> Description of Artificial Sequence: synthetic DNA
 <400> 21
 ggtggatctg ggacagatta tactctcacc atcagcagcc tgcagcctga agattttgca 60
 acttattact gtcacagta tagtaagctt cc 92
 45

<210> 22
 <211> 80
 <212> DNA
 50 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 22
 55 gttttcccag tcacgaccgt acgtttaatc tctaccttgg tccctggcc gaacgtccac 60

ggaagcttac tatactgatg 80

(0232)

5 <210> 23
 <211> 45
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 23
 agcttccatg gacgttcggt ggaggcacca agctggaaat caaac 45

15 <210> 24
 <211> 45
 <212> DNA
 <213> Artificial Sequence

20 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 24
 gtacgtttga ttccagett ggtgcctcca ccgaacgtcc atgga 45

25 <210> 25
 <211> 29
 <212> DNA
 <213> Artificial Sequence

30 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 25
 gatgcaggca gggaggatgc agtctgggt 29

35 <210> 26
 <211> 29
 <212> DNA
 <213> Artificial Sequence

40 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 26
 acccagactg catcctccct gcctgcac 29

45 (0233)

50 <210> 27
 <211> 384
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <220>

55 <221> CDS

<222> (1).. (384)

<400> 27

atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48

Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Leu Cys Phe Gln

5 -20 -15 -10 -5

ggt acc aga tgt gac atc cag atg acc cag act gca tcc tcc ctg cct 96

Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Thr Ala Ser Ser Leu Pro

-1 1 5 10

10

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144

Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp

15 20 25

15

att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc cct 192

Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro

30 35 40

20

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240

Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser

45 50 55 60

25

agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288

Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser

65 70 75

30

agc ctg cag cct gaag gat ttt gca act tat tac tgt cat cag tat agt 336

Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys His Gln Tyr Ser

80 85 90

aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384

Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg

95 100 105

35

(0234)

<210> 28

<211> 26

40

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: synthetic DNA

<400> 28

45

actgatgaca gaaataagtt gcaaaa

26

<210> 29

<211> 26

50

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: synthetic DNA

<400> 29

55

ttttgcaact tatttctgtc atcagt

26

<210> 30
 <211> 384
 5 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <220>
 10 <221> CDS
 <222> (1)..(384)
 <400> 30
 atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
 Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
 15 -20 -15 -10 -5
 ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96
 Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
 -1 1 5 10
 20 gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
 Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
 15 20 25
 25 att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc cct 192
 Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro
 30 35 40
 aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240
 Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
 45 50 55 60
 agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288
 Arg Phe Ser Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser
 35 65 70 75
 agc ctg cag cct gaa gat ttt gca act tat ttc tgt cat cag tat agt 336
 Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Phe Cys His Gln Tyr Ser
 80 85 90
 40 aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384
 Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
 95 100 105
 45 (0235)
 <210> 31
 <211> 28
 50 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 31
 55 ggagcttaac ggctttgtct ggtttctg

28

<210> 32
 <211> 28
 5 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 <400> 32
 10 cagaaaccag acbaagccgt taagctcc 28

(0236)

15 <210> 33
 <211> 384
 <212> DNA
 <213> Artificial Sequence
 <220>
 20 <223> Description of Artificial Sequence: synthetic DNA
 <220>
 <221> CDS
 <222> (1)..(384)
 <400> 33

25 atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
 Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
 -20 -15 -10 -5

30 ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96
 Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
 -1 1 5 10

35 gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
 Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
 15 20 25

40 att agt aat tat tta aac tgg tat cag cag aaa cca gac aaa gcc gtt 192
 Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Asp Lys Ala Val
 30 35 40

45 aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240
 Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
 45 50 55 60

50 agc ctg cag cct gaa gat ttt gca act tat tac tgt cat cag tat agt 336
 Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys His Gln Tyr Ser
 80 85 90

55 aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384
 Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
 95 100 105

(0237)

5 <210> 34
<211> 48
<212> DNA
<213> Artificial Sequence
<220>
10 <223> Description of Artificial Sequence: synthetic DNA
<400> 34
gttgcgatat cttcaggctg cagattgctg atggtgagac tataatct 48

15 <210> 35
<211> 48
<212> DNA
<213> Artificial Sequence
<220>
20 <223> Description of Artificial Sequence: synthetic DNA
<400> 35
agattatagt ctcaccatca gcaatctgca gccgtgaagat atcgcaac 48

25 <210> 36
<211> 384
<212> DNA
<213> Artificial Sequence
<220>
30 <223> Description of Artificial Sequence: synthetic DNA
<220>
<221> CDS
<222> (1)..(384)
<400> 36

35 atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
-20 -15 -10 -5

40 ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96
Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
-1 1 5 10

45 gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
15 20 25

50 att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc cct 192
Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro
30 35 40

55 aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240
Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
45 50 55 60

agg ttc agc ggc ggt gga tct ggg aca gat tat agt ctc acc atc agc 288

Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser
65 70 75

5 aat ctg cag cct gaa gat atc gca act tat tac tgt cat cag tat agt 336
Asn Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys His Gln Tyr Ser
80 85 90

10 aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384
Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
95 100 105

(0238)

15 <210> 37
<211> 25
<212> DNA
<213> Artificial Sequence
<220>
20 <223> Description of Artificial Sequence: synthetic DNA
<400> 37
ttcaggtgc agattgctga tgggtg 25

25 <210> 38
<211> 25
<212> DNA
<213> Artificial Sequence
<220>
30 <223> Description of Artificial Sequence: synthetic DNA
<400> 38
caccatcagc aatctgcagc ctgaa 25

35 <210> 39
<211> 384
<212> DNA
<213> Artificial Sequence
<220>
40 <223> Description of Artificial Sequence: synthetic DNA
<220>
<221> CDS
<222> (1)..(384)
<400> 39
45 atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln
-20 -15 -10 -5

50 ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96
Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
-1 1 5 10

55 gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
15 20 25

att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc cct 192
 Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro
 30 35 40

5 aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240
 Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
 45 50 55 60

10 agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288
 Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser
 65 70 75

15 aat ctg cag cct gaa gat ttt gca act tat tac tgt cat cag tat agt 336
 Asn Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys His Gln Tyr Ser
 80 85 90

20 aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384
 Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
 95 100 105

(0239)

25 <210> 40
 <211> 30
 <212> DNA
 <213> Artificial Sequence
 <220>

30 <223> Description of Artificial Sequence: synthetic DNA
 <400> 40
 gacagaaata agttgcgata tcttcaggct 30

35 <210> 41
 <211> 30
 <212> DNA
 <213> Artificial Sequence
 <220>

40 <223> Description of Artificial Sequence: synthetic DNA
 <400> 41
 agcctgaaga tatcgcaact tattctgtc 30

45 <210> 42
 <211> 384
 <212> DNA
 <213> Artificial Sequence
 <220>

50 <223> Description of Artificial Sequence: synthetic DNA
 <220>
 <221> CDS
 <222> (1)..(384)
 <400> 42

55 atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48

Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Leu Cys Phe Gln
-20 -15 -10 -5

5 ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96
Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
-1 1 5 10

10 gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
15 20 25

15 att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc cct 192
Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro
30 35 40

20 aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240
Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
45 50 55 60

25 agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288
Arg Phe Ser Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser
65 70 75

30 agc ctg cag cct gaa gat atc gca act tat ttc tgt cat cag tat agt 336
Ser Leu Gln Pro Glu Asp Ile Ala Thr Tyr Phe Cys His Gln Tyr Ser
80 85 90

35 aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384
Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
95 100 105

(0240)

35 <210> 43
<211> 87
<212> DNA
<213> Artificial Sequence
<220>

40 <223> Description of Artificial Sequence: synthetic DNA
<400> 43
gcactacaag tgatggtagac tctgtctcct acagatgcag gcaggaggga tgcagactgg 60
gtcatctgga tgacacatct ggtacct 87

45

50 <210> 44
<211> 89
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: synthetic DNA
<400> 44
gtcaccatca cttgtagtgc aagtcaggac attagtaatt atttaaacctg gtatcagcag 60

55

aaaccaggga aagccgttaa gtcctgat

89

<210> 45

5 <211> 89

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: synthetic DNA

10 <400> 45

taatctgtcc cagatccacc gccgctgaac ctgatggga ccccgagtg taaatttgat 60

gagtaaaaga tcaggagctt aacggcttt

89

15

<210> 46

<211> 92

<212> DNA

<213> Artificial Sequence

20 <220>

<223> Description of Artificial Sequence: synthetic DNA

<400> 46

ggtagatctg ggacagatta tactctcacc atcagcagcc tgcagcctga agattttgca 60

25 acttatttct gtcacagta tagtaagctt cc

92

(0241)

30 <210> 47

<211> 384

<212> DNA

<213> Artificial Sequence

<220>

35 <223> Description of Artificial Sequence: synthetic DNA

<220>

<221> CDS

<222> (1)..(384)

<400> 47

40 atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48

Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Cys Phe Gln

-20 -15 -10 -5

ggc acc aga tgt gac atc cag atg acc cag tct gca tcc tcc ctg cct 96

45 Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Ala Ser Ser Leu Pro

-1 1 5 10

gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144

50 Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp

15 20 25

att agt aat tat tta aac tgg tat cag cag aaa cca ggg aaa gcc gtt 192

Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Val

30 35 40

55

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240
 Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
 45 50 55 60

5 agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288
 Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser
 65 70 75

10 agc ctg cag cct gaa gat ttt gca act tat ttc tgt cat cag tat agt 336
 Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Phe Cys His Gln Tyr Ser
 80 85 90

15 aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384
 Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
 95 100 105

(0242)

20 <210> 48
 <211> 384
 <212> DNA
 <213> Artificial Sequence
 <220>
 25 <223> Description of Artificial Sequence: synthetic DNA
 <220>
 <221> CDS
 <222> (1)..(384)
 <400> 48

30 atg atg tcc tct gct cag ttc ctt ggt ctc ctg ttg ctc tgt ttt caa 48
 Met Met Ser Ser Ala Gln Phe Leu Gly Leu Leu Leu Leu Cys Phe Gln
 -20 -15 -10 -5

35 ggt acc aga tgt gac atc cag atg acc cag tct cca tcc tcc ctg tct 96
 Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
 -1 1 5 10

40 gca tct gta gga gac aga gtc acc atc act tgt agt gca agt cag gac 144
 Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp
 15 20 25

45 att agt aat tat tta aac tgg tat cag cag aaa cca gac aaa gcc gtt 192
 Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Asp Lys Ala Val
 30 35 40

aag ctc ctg atc ttt tac tca tca aat tta cac tcg ggg gtc cca tca 240
 Lys Leu Leu Ile Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser
 45 50 55 60

50 agg ttc agc ggc ggt gga tct ggg aca gat tat act ctc acc atc agc 288
 Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser
 65 70 75

55 agc ctg cag cct gaa gat atc gca act tat ttc tgt cat cag tat agt 336
 Ser Leu Gln Pro Glu Asp Ile Ala Thr Tyr Phe Cys His Gln Tyr Ser

	80	85	90
	aag ctt ccg tgg acg ttc ggc cag ggg acc aag gta gag att aaa cgt 384		
5	Lys Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg		
	95	100	105
	(0243)		
10	<210> 49		
	<211> 76		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
15	<223> Description of Artificial Sequence: synthetic DNA		
	<400> 49		
	catgcatgag gctctgcaca accactacac gcagaagagc ctctccctgt ctcccggggg 60		
20	agaattcatt gatcag 76		
	<210> 50		
	<211> 85		
	<212> DNA		
25	<213> Artificial Sequence		
	<220>		
	<223> Description of Artificial Sequence: synthetic DNA		
	<400> 50		
	gatccctgac aatgaattct cccccgggag acagggagag gctcttctgc gtgtagtggt 60		
30	tgtgcagagc ctcatgcatg gggcc 85		
	<210> 51		
35	<211> 37		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Description of Artificial Sequence: synthetic DNA		
40	<400> 51		
	gtctcccggg aaagcaccta ctagtagttc tacaag 37		
	<210> 52		
45	<211> 38		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Description of Artificial Sequence: synthetic DNA		
50	<400> 52		
	ccctgatcaa tgaattcaag tcagtgttga gatgatgc 38		
	(0244)		
55			

<210> 53
 <211> 582
 <212> PRT
 <213> Artificial Sequence

5 <400> 53
 Glu Val Gln Leu Val Glu Ser Gly Gly Asp Phe Val Gln Pro Gly Gly
 1 5 10 15
 10 Ser Leu Arg Val Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser His Tyr
 20 25 30
 Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 15 Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 20 Leu Gln Met Arg Ser Leu Arg Ala Glu Asp Ser Ala Val Tyr Phe Cys
 85 90 95
 25 Thr Arg Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser Trp Gly Gln Gly
 100 105 110
 Thr Leu Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
 115 120 125
 30 Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
 130 135 140
 Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
 145 150 155 160
 35 Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
 165 170 175
 40 Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
 180 185 190
 Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro
 195 200 205
 45 Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys
 210 215 220
 Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro
 225 230 235 240
 50 Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser
 245 250 255
 55 Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp
 260 265 270

	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	
		275						280					285				
5	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	
		290					295					300					
	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	
	305					310					315					320	
10	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	
					325					330					335		
	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	
15				340					345					350			
	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	
			355					360					365				
20	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	
	370						375					380					
	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	
	385					390					395					400	
25	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	
				405					410						415		
	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	
30				420					425					430			
	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	
		435						440					445				
35	Lys	Ala	Pro	Thr	Ser	Ser	Ser	Thr	Lys	Lys	Thr	Gln	Leu	Gln	Leu	Glu	
	450					455						460					
	His	Leu	Leu	Leu	Asp	Leu	Gln	Met	Ile	Leu	Asn	Gly	Ile	Asn	Asn	Tyr	
	465					470					475					480	
40	Lys	Asn	Pro	Lys	Leu	Thr	Arg	Met	Leu	Thr	Phe	Lys	Phe	Tyr	Met	Pro	
				485					490					495			
	Lys	Lys	Ala	Thr	Glu	Leu	Lys	His	Leu	Gln	Cys	Leu	Glu	Glu	Glu	Leu	
45			500						505					510			
	Lys	Pro	Leu	Glu	Glu	Val	Leu	Asn	Leu	Ala	Gln	Ser	Lys	Asn	Phe	His	
		515					520						525				
50	Leu	Arg	Pro	Arg	Asp	Leu	Ile	Ser	Asn	Ile	Asn	Val	Ile	Val	Leu	Glu	
	530					535						540					
	Leu	Lys	Gly	Ser	Glu	Thr	Thr	Phe	Met	Cys	Glu	Tyr	Ala	Asp	Glu	Thr	
	545					550					555					560	
55																	

Ala Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser
565 570 575

5 Ile Ile Ser Thr Leu Thr
580

(0245)

10 <210> 54
<211> 108
<212> PRT
<213> Artificial Sequence
<400> 54

15 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

20 Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr
20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Asp Lys Ala Val Lys Leu Leu Ile
35 40 45

25 Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Gly Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

30 Glu Asp Ile Ala Thr Tyr Phe Cys His Gln Tyr Ser Lys Leu Pro Trp
85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
100 105

35

<210> 55
<211> 119
<212> PRT
40 <213> Mus musculus
<400> 55

Glu Val Thr Leu Val Glu Ser Gly Gly Asp Phe Val Lys Pro Gly Gly
1 5 10 15

45 Ser Leu Lys Val Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser His Tyr
20 25 30

Ala Met Ser Trp Val Arg Gln Thr Pro Ala Lys Arg Leu Glu Trp Val
35 40 45

50

Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val
50 55 60

55 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Arg Ser Leu Arg Ser Glu Asp Ser Ala Met Tyr Phe Cys
 85 90 95
 5 Thr Arg Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser Trp Gly Gln Gly
 100 105 110
 Thr Thr Leu Thr Val Ser Ser
 115
 10 (0246)
 <210> 56
 <211> 108
 15 <212> PRT
 <213> Mus musculus
 <400> 56
 Asp Ile Gln Met Thr Gln Thr Ala Ser Ser Leu Pro Ala Ser Leu Gly
 1 5 10 15
 20 Asp Arg Val Thr Ile Ser Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr
 20 25 30
 Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val Lys Leu Leu Ile
 25 35 40 45
 Phe Tyr Ser Ser Asn Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 30 Gly Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser Asn Leu Glu Pro
 65 70 75 80
 Glu Asp Ile Ala Thr Tyr Phe Cys His Gln Tyr Ser Lys Leu Pro Trp
 85 90 95
 35 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
 100 105
 40 <210> 57
 <211> 582
 <212> PRT
 <213> Artificial Sequence
 <400> 57
 45 Glu Val Thr Leu Val Glu Ser Gly Gly Asp Phe Val Lys Pro Gly Gly
 1 5 10 15
 Ser Leu Lys Val Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser His Tyr
 20 25 30
 50 Ala Met Ser Trp Val Arg Gln Thr Pro Ala Lys Arg Leu Glu Trp Val
 35 40 45
 Ala Tyr Ile Ser Ser Gly Gly Ser Gly Thr Tyr Tyr Ser Asp Ser Val
 55 50 55 60

	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr	65	70	75	80
5	Leu Gln Met Arg Ser Leu Arg Ser Glu Asp Ser Ala Met Tyr Phe Cys	85	90	95	
	Thr Arg Val Lys Leu Gly Thr Tyr Tyr Phe Asp Ser Trp Gly Gln Gly	100	105	110	
10	Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe	115	120	125	
	Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu	130	135	140	
15	Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp	145	150	155	160
20	Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu	165	170	175	
	Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser	180	185	190	
25	Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro	195	200	205	
	Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys	210	215	220	
30	Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro	225	230	235	240
35	Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser	245	250	255	
	Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp	260	265	270	
40	Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn	275	280	285	
45	Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val	290	295	300	
	Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu	305	310	315	320
50	Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys	325	330	335	
	Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr	340	345	350	
55					

Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr
 355 360 365
 5 Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu
 370 375 380
 Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu
 385 390 395 400
 10 Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys
 405 410 415
 Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
 420 425 430
 15 Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 435 440 445
 Lys Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu
 20 450 455 460
 His Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr
 465 470 475 480
 25 Lys Asn Pro Lys Leu Thr Arg Met Leu Thr Phe Lys Phe Tyr Met Pro
 485 490 495
 Lys Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu
 30 500 505 510
 Lys Pro Leu Glu Glu Val Leu Asn Leu Ala Gln Ser Lys Asn Phe His
 515 520 525
 Leu Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu
 35 530 535 540
 Leu Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr
 545 550 555 560
 40 Ala Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser
 565 570 575
 Ile Ile Ser Thr Leu Thr
 580
 45

(Brief description of the drawings)

- (Fig. 1) is a drawing showing construction steps of plasmid phKM641H.
 (Fig. 2) is a drawing showing construction steps of plasmid phKM641L.
 (Fig. 3) is a drawing showing construction steps of plasmid pT796H641L.
 50 (Fig. 4) is a drawing showing construction steps of plasmid pBS641H.
 (Fig. 5) is a drawing showing construction steps of plasmid pT641.

(Fig. 6) is a drawing showing construction steps of plasmid pT641HCDR.

(Fig. 7) is a drawing showing construction steps of plasmids pT641LCDR and pT641HLCDR.

(Fig. 8) is a drawing showing activity evaluation of an anti-GD3 chimeric antibody and an anti-GD3 CDR-grafted antibody by their transient expression using plasmids pT641, pT641HCDR, pT641LCDR and pT641HLCDR. The ordinate and the abscissa represent the expression vector name and the relative activity (%) when the activity of anti-GD3 chimeric antibody is defined as 100, respectively.

(Fig. 9) is a drawing showing construction steps of plasmid phKM641Lm-69.

(Fig. 10) is a drawing showing construction steps of plasmids pT641HLCDRNL, pT641HLCDRm-1, pT641HLCDRm-4, pT641HLCDRm-6, pT641HLCDRm-7, pT641HLCDRm-8, pT641HLCDRm-9 and pT641HLCDRm-69.

(Fig. 11) is a drawing showing activity evaluation of anti-GD3 chimeric antibody and anti-GD3 CDR-grafted antibody by their transient expression using plasmids pT641, pT641HLCDR, pT641HLCDRNL, pT641HLCDRm-1, pT641HLCDRm-4, pT641HLCDRm-6, pT641HLCDRm-7, pT641HLCDRm-8, pT641HLCDRm-9 and pT641HLCDRm-69. The ordinate and the abscissa represent the expression vector name and the relative activity (%) when the activity of anti-GD3 chimeric antibody is defined as 100, respectively.

(Fig. 12) is a drawing showing construction steps of plasmids pKANTEX641HLCDR, pKANTEX641HLCDRm-9 and pKANTEX641HLCDRm-69.

(Fig. 13) is a drawing showing SDS-PAGE (using a 4 to 15% gradient gel) electrophoresis patterns of purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871. The left side shows results of electrophoresis carried out under non-reducing conditions, and the right side shows those under reducing conditions. Lanes 1, 2, 3, 4, 5, 6, 7 and 8 show electrophoresis patterns of high molecular weight markers, KM8869, KM8870, KM8871, low molecular weight markers, KM8869, KM8870 and KM8871, respectively.

(Fig. 14) is a drawing showing binding activity of purified anti-GD3 chimeric antibody KM871 and purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 to GD3 measured by changing the antibody concentration. The ordinate and the abscissa are the binding activity to GD3 and the antibody concentration,

respectively. "○", "○", "□" and "□" show the activities of KM871, KM8869, KM8870 and KM8871, respectively.

(Fig. 15) is a drawing showing binding activity of purified anti-GD3 chimeric antibody KM871 and purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 to GD3 measured by changing an amount of GD3 to be adsorbed to a plate. The ordinate and the abscissa are the binding activity to GD3 and the amount of GD3 adsorbed to the plate, respectively. "○", "○", "□" and "□" show the activities of KM871, KM8869, KM8870 and KM8871, respectively.

(Fig. 16) is a drawing showing reactivity of purified anti-GD3 chimeric antibody KM871 and purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 with various gangliosides. The ordinate and the abscissa are the kind of ganglioside and the binding activity, respectively. AcGM2, GcGM2, AcGM3 and GcGM3 indicate N-acetylGM2, N-glycolylGM2, N-acetylGM3 and N-glycolylGM3. "□", "□", "□" and "□" show the reactivities of KM871, KM8869, KM8870 and KM8871, respectively.

(Fig. 17) is a drawing showing reactivity of purified anti-GD3 chimeric antibody KM871 and purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 with human melanoma cell lines G-361 and SK-MEL-28. The ordinate and the abscissa are the number of cells and the fluorescence intensity, respectively. The graphs show reactivities of control, KM871, KM8869, KM8870 and KM8871, respectively, from the bottom.

(Fig. 18) is a drawing showing CDC activity of purified anti-GD3 chimeric antibody KM871 and purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 against human melanoma cell lines G-361 and SK-MEL-28. The ordinate and the abscissa are the cytotoxic activity and the antibody concentration, respectively. "□", "□", "□" and "□" show the reactivities of KM871, KM8869, KM8870 and KM8871, respectively.

(Fig. 19) is a drawing showing ADCC activity of purified anti-GD3 chimeric antibody KM871 and purified anti-GD3 CDR-grafted antibodies KM8869, KM8870 and KM8871 against human melanoma cell lines G-361 and SK-MEL-28. The ordinate and the abscissa are the cytotoxic activity and the antibody concentration, respectively. "□", "□", "□" and "□" show the reactivities of KM871, KM8869, KM8870 and KM8871, respectively.

(Fig. 20) is a drawing showing construction steps of plasmid pBSA-B.

(Fig. 21) is a drawing showing construction steps of plasmid pBSΔhCyl-IL-

2.

(Fig. 22) is a drawing showing construction steps of plasmid pBShCyl-IL-2.

(Fig. 23) is a drawing showing construction steps of plasmid pKANTEX8871-hIL-2.

(Fig. 24) is a drawing showing SDS-PAGE (using a 4 to 15% gradient gel) electrophoresis patterns of purified anti-GD3 CDR-grafted antibody KM8871 and purified fusion protein KM8871-hIL-2. The left side and the right side are results of electrophoresis carried out under non-reducing conditions, and those under reducing conditions, respectively. Lanes 1, 2, 3, 4, 5 and 6 show electrophoresis patterns of low molecular weight markers, KM8871, KM8871-hIL-2, low molecular weight markers, KM8871 and KM8871-hIL-2, respectively.

(Fig. 25) is a drawing showing binding activity of purified anti-GD3 CDR-grafted antibody KM8871 and purified fusion protein KM8871-hIL-2 to GD3 measured by changing the antibody concentration. The ordinate and the abscissa are the binding activity to GD3 and the antibody concentration. "○" and "○" show the activities of KM8871 and of KM8871-hIL-2, respectively.

(Fig. 26) is a drawing showing reactivity of purified anti-GD3 CDR-grafted antibody KM8871 and purified fusion protein KM8871-hIL-2 with various gangliosides. The ordinate and the abscissa are the kind of ganglioside and the binding activity. AcGM2, GcGM2, AcGM3 and GcGM3 indicate N-acetylGM2, N-glycolylGM2, N-acetylGM3 and N-glycolylGM3, respectively. "□" and "□" show the reactivities of KM8871 and KM8871-hIL-2, respectively.

(Fig. 27) is a drawing showing reactivity of purified anti-GD3 CDR-grafted antibody KM8871 and purified fusion protein KM8871-hIL-2 with human melanoma cell line G-361. The ordinate and the abscissa are the number of cells and the fluorescence intensity, respectively. The drawings show the reactivities of KM8871 and KM8871-hIL-2, respectively, from the upper column.

(Fig. 28) is a drawing showing growth-supporting activity of hIL-2 and purified fusion protein KM8871-hIL-2 against hIL-1-dependent cell CTLL-2 measured by changing concentration of each protein. The ordinate and the abscissa are the growth-supporting activity and the protein concentration, respectively. "○" and "○" indicate the activities of hIL-2 and KM8871-hIL-2, respectively.

(Fig. 29) is a drawing showing results of the measurement of the activation and accompanying cytotoxic activity of human effector cells by purified anti-GD3 CDR-grafted antibody KM8871 and purified fusion protein KM8871-hIL-2. The ordinate and the abscissa are the cytotoxic activity and the used protein, respectively. "□" and "□" indicate the activities of KM8871 and KM8871-hIL-2, respectively.

(Fig. 30) is a drawing showing construction steps of plasmid pKANTEX871-hIL-2.

(Fig. 31) is a drawing showing SDS-PAGE (using a 4 to 15% gradient gel) electrophoresis patterns of anti-GD3 chimeric antibody KM871 and purified fusion protein KM871-hIL-2. The left side and the right side are results of electrophoresis carried out under reducing conditions and under non-reducing conditions, respectively. Lanes 1, 2, 3, 4, 5 and 6 show electrophoresis patterns of low molecular weight markers, KM871, KM871-hIL-2, low molecular weight markers, KM871 and KM871-hIL-2, respectively.

(Fig. 32) is a drawing showing binding activity of anti-GD3 chimeric antibody KM871 and purified fusion protein KM871-hIL-2 to GD3 measured by changing the antibody concentration. The ordinate and the abscissa are the binding activity to GD3 and the antibody concentration, respectively. "○" and "□" show the activities of KM871 and KM871-hIL-2, respectively.

(Fig. 33) is a drawing showing reactivity of anti-GD3 chimeric antibody KM871 and purified fusion protein KM871-hIL-2 with various gangliosides. The ordinate and the abscissa are the kind of ganglioside and the binding activity, respectively. AcGM2, GcGM2, AcGM3 and GcGM3 indicate N-acetylGM2, N-glycolylGM2, N-acetylGM3 and N-glycolylGM3, respectively. "□" and "□" show the reactivities of KM871 and KM871-hIL-2, respectively.

(Fig. 34) is a drawing showing reactivity of anti-GD3 chimeric antibody KM871 and purified fusion protein KM871-hIL-2 with human melanoma cell lines G-361 and SK-MEL-28. The ordinate and the abscissa are the number of cells and the fluorescence intensity, respectively. The graphs show the reactivities of KM871 to G361 cell, KM871-hIL-2 to G361 cell, KM871 to SK-MEL-28 and KM871-hIL-2 to SK-MEL-28, respectively, from the upper column.

(Fig. 35) is a drawing showing growth-supporting activity of hIL-2 and purified fusion protein KM871-hIL-2 against hIL-2-dependent cell CTLL-2 measured by changing concentration of each protein. The ordinate and the abscissa are the growth-supporting activity and the protein concentration, respectively. "○" and "○" indicate the activities of hIL-2 and KM871-hIL-2, respectively.

(Fig. 36) is a drawing showing results of the measurement of the activation and accompanying cytotoxic activity of human effector cells by anti-GD3 chimeric antibody KM871 and purified fusion protein KM871-hIL-2. The ordinate and the abscissa are the cytotoxic activity and the used protein, respectively. "□" and "□" indicate activities of KM871 and KM871-hIL-2, respectively.

(Fig. 37) is a drawing showing construction steps of plasmid pKANTEXGD3.

(Fig. 38) is a drawing showing reactivity of anti-GD3 chimeric antibody KM871 with GD3 synthase gene-introduced B16 cell lines B16-29-10 and B16-8-3 and the parent line B16-F10 cell. The ordinate and the abscissa are the number of cells and the fluorescence intensity, respectively. The drawings show the reactivities of B16-F10 cell, B16-29-10 and B16-8-3, respectively, from the upper column.

(Fig. 39) is a drawing showing results of the measurement of anti-metastatic effects of anti-GD3 chimeric antibody KM871 and KM871-hIL-2 on BL/6 mice into which a GD3-positive B16 cell line B16-29-10 is transplanted. The ordinate and the bar of the graph show the number of metastatic foci on the lung surface and the average value of each group, respectively. No administration means a control group in which the antibody is not administered. "◇", "○" and "○" indicate the numbers of lung-metastasized foci in mice of the control group, the KM871-hIL-2-administered group and the KM871-administered group, respectively.

(Fig. 40) is a drawing showing a result of the measurement of anti-metastatic effects of anti-GD3 chimeric antibody KM871 and KM871-hIL-2 on BL/6 mice into which a GD3-positive B16 cell line B16-8-3 is transplanted. The ordinate and the bar of the graph show the number of metastatic foci on the lung surface and the average value of each group, respectively. No administration means a control group in which the antibody is not administered. "◇", "○" and "○" indicate the numbers of lung-metastasized foci in mice of the control group, the KM871-hIL-2-administered group and the KM871-administered group, respectively.

(Fig. 41) is a drawing showing results of the measurement of anti-metastatic effects by combined administration of anti-GD3 chimeric antibody KM871 and hIL-2 and administration of KM871-hIL-2 on BL/6 mice into which a GD3-positive B16 cell line B16-29-10 is transplanted. The ordinate and the bar of the graph show the number of metastatic foci on the lung surface and the average value of each group. No administration means a control group in which the antibody is not administered. "◇", "○" and "○" indicate the number of lung-metastasized foci in mice of the control group, the KM871-hIL-2-administered group and the KM871/hIL-2 combined administered group, respectively.

(Document name)

Drawings

FIG.1




FIG.2




FIG.3




FIG.4




FIG.5




FIG.6




FIG.7




FIG.8




FIG.9




FIG.10




FIG.11




FIG.12




FIG.13




FIG.14




FIG.15




FIG.16




FIG.17




FIG.18





FIG.19





FIG.20




FIG.21




FIG.22




FIG.23




FIG.24




FIG.25




FIG.26





FIG.27



G361CELL




FIG.28




FIG.29




FIG.30




FIG.31




FIG.32




FIG.33




FIG.34







FIG.35




FIG.36




FIG.37




FIG.38






FIG.39




FIG.40





FIG.41



(Document name)

Abstract

(Abstract)

(Problems)

5 If derivatives of a human CDR-grafted antibody or an anti-GD3 antibody which specifically with tumor cells with cytokine or the like are produced, they are expected to be used as new diagnostic or therapeutic agents for human cancer.

(Means for solution)

10 The present invention relates to a human CDR-grafted antibody against ganlioside GD3 (hereinafter referred to "GD3"), derivatives of an anti-GD3 antibody and cytokine, and use for treatment and diagnosis of the antibody and the derivatives.

(Selected drawings)

None

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.